

The copyright of this thesis vests in the author. No quotation from it or information derived from it is to be published without full acknowledgement of the source. The thesis is to be used for private study or non-commercial research purposes only.

Published by the University of Cape Town (UCT) in terms of the non-exclusive license granted to UCT by the author.

Keeping time on the plant-pathogen arms race: A role for the plant circadian clock in immune response

By Vaibhav Bhardwaj

Dissertation presented for the degree of Master of Science in the
Department of Molecular and Cellular Biology, Faculty of Science at
the University of Cape Town

August 2011

Plagiarism declaration

I know the meaning of plagiarism and declare that all of the work in this dissertation, save for that which is properly acknowledged is my own.

Vaibhav Bhardwaj

University of Cape Town

Abstract

Plants and pathogens have been engaged in an evolutionary arms race that has led to both species evolving a complex set of defence mechanisms. Initially, plants perceive invading bacteria by recognising microbe associated molecular patterns (MAMPs). This leads to the MAMP triggered immune response (MTI). Bacteria in turn evolved effector proteins that act to repress MTI leading to virulence. However, plant resistance (R) proteins, can recognise these effectors, activating effector-triggered immunity (ETI). While plants are capable of initiating extreme defence responses such as programmed cell death, they display a carefully regulated defence response. Recently, investigations of defence responses to the oomycete *Hyaloperonospora arabidopsis* revealed a molecular link between a circadian clock component and plant defence. A systems approach revealed that a number of defence genes showed peak expression at dawn and possessed CIRCADIAN CLOCK ASSOCIATED1 (CCA1) binding sites in their promoters. In this study, *Arabidopsis thaliana* (*Arabidopsis*) in the Columbia-0 (Col-0) background showed time-of-day variation in susceptibility to the plant-pathogen *Pseudomonas syringae* DC3000 pathovar tomato (*P. syringae* DC3000) when infected under constant light and temperature conditions. Wild type plants showed least susceptibility at circadian time (CT) 26 and 50, which correspond to 'subjective' morning. Plants were most susceptible when infected at CT42 and CT66, 'subjective' night. Additionally, the arrhythmic *CCA1*-over-expressor (*CCA1-ox*) line and *elf3-1* mutant revealed similar levels of susceptibility at both subjective morning and night infections. Callose deposition is an early MAMP defence response. Measuring callose levels in response to infection with non-pathogenic *P. syringae* *hrpA* demonstrated that callose production also showed time-of-day variation. Higher callose levels were observed after infection at CT26 compared to CT42 in wild type plants, while *CCA1-ox* showed similar levels after infection, at both time points. This result complemented the earlier observation. Paradoxically, promoter activity of *Pathogenesis related1* (*PR1*) a defence gene, indicated quicker induction at CT42 than at CT26 while promoter activity of *Oxidative Stress Induced 1* (*OXI1*) showed no time-of-day variation in activity. Preliminary gene expression measurements indicated that induction of *WRKY29* showed time-of-day variation with greater induction at CT26 than at CT42 in wild type plants but *WRKY22* did not. Analysis of basal expression levels of a gene encoding a suppressor of R proteins (*SRFR1*) also showed time of day variation with higher levels at CT42. While clock control in plant defence is important, measuring promoter activity of *COLD AND CIRCADIAN REGULATED 2* (*CCR2*), a clock controlled output gene, revealed core circadian clock function is unaffected by pathogen infection. Collectively these results indicate that there are clock co-ordinated defence responses in *Arabidopsis* and that plant defence against *P. syringae* is greatest at dawn.

Acknowledgements

During my M.Sc in the Molecular and Cellular Biology department at UCT, my supervisors, Dr. Laura Roden and Dr. Robert Ingle, have been continually supportive of all my efforts in the lab.

I would also like to thank Mariette Smart who guided me through my early stages in the lab and for being a role model in and out of the lab. Her friendship and support made every day in the lab worth it.

My sincere gratitude to Alykhan Vira for the unlimited time he offered to me to discuss various ideas that helped with my experiments and my writing. I also thank Ragesh Pillai for his support and friendship and for all the lifts back from the labs at 2 am.

Many thanks to my lab colleagues, Lovemore Kunorozva, Eric Banda, Stewart Norman and Fatime Ramla who helped with some of the experiments.

My sincere gratitude for the support of the Mellon Mays Foundation and Kathy Erasmus for the opportunities to travel twice to conferences in Chicago and Princeton University where I had the opportunity to discuss my work with renowned academics.

Thank you also to the two most important women in my life, my mother and my high school biology teacher, Mrs Atkinson for always believing in me and for being amazing role models.

Lastly, thank you to members of the MCB department of UCT who were a constant source of motivation.

Contents

Plagiarism declaration.....	i
Abstract.....	ii
Acknowledgements.....	iii
List of Figures	vi
List of tables:.....	vii
Abbreviations	viii
Chapter 1: Literature Review	1
The <i>Arabidopsis</i> immune system	1
MAMP triggered immune response (MTI)	2
Effector Triggered Immune response (ETI)	5
The two-branched paradigm challenged:	6
Negative regulation of plant defence	8
Comparing innate immunity in plants, animals, insects and invertebrates:	11
The Circadian Clock and Immunity	12
The Circadian Clock of <i>Drosophila melanogaster</i> and its role in immunity.....	13
Circadian regulation of immune responses in mammals	15
The Plant Circadian Clock and its role in Immunity	16
Concluding remarks	21
Research question.....	21
Chapter 2: Materials and Methods.....	23
2.1 Plant Materials	23
2.2 Plant growth conditions.....	23
2.3 Bacterial growth conditions.....	24
2.4 Pressure infiltrating <i>Arabidopsis</i> plants for pathogen assays.....	25
2.5 Callose deposition assays.....	25
2.6 Measuring gene expression	25
2.6.1 Gene expression using luciferase.....	25
2.6.2 Quantitative PCR	26
2.7 Online Bioinformatics tools.....	28
2.8 Statistical analysis	28
Chapter 3: Results	29
3.1 Investigating the MAMP-triggered immune response for circadian control.....	29
3.1.1 Investigating the effect of time-of-day on <i>Arabidopsis</i> response to <i>P. syringae</i> DC3000: ..	29

3.1.2 Investigating <i>elf3-1</i> and <i>CCA1-ox</i> for time-of-day variation in susceptibility to <i>P. syringae</i> DC3000:.....	30
3.1.3 Investigating time-of-day variation in <i>CCA1-ox</i> in the second cycle:.....	32
3.2 Investigating the ETI response of <i>Arabidopsis</i> for time-of-day variation under free running conditions:	34
3.2.1 Infecting wild type and clock mutants with <i>P. syringae</i> DC3000 <i>avrB</i> and <i>avrRpt2</i> at CT26 and CT42:	34
3.3 Measuring callose levels of <i>Arabidopsis</i> in response to <i>P. syringae hrpA</i> at CT26 and CT42:	35
3.4 Promoter activity and expression of genes involved in the MTI response to <i>P. syringae</i> DC3000 infection:	37
3.4.1 Promoter activity of <i>OXI1</i> following infection	37
3.4.2 Promoter activity of <i>PR1</i> following infection	38
3.4.3 Measuring transcript levels of <i>WRKY</i> transcription factors	39
3.4.4 <i>SRFR1</i> levels at CT26 and CT42	41
3.4.5 Investigating circadian clock function during infection	42
Chapter 4 Discussion.....	44
Concluding remarks	51
References	52

List of Figures

Figure 1. An illustration of the EMT and SA sectors showing antagonistic associations- The 'two-hit' model of plant defence.....	9
Figure 2. An illustration explaining circadian time and free-running conditions that the plants were subjected to..	24
Figure 3.1 <i>Arabidopsis thaliana</i> (Col-0) shows variable susceptibility to <i>P. syringae</i> DC3000 under free-running conditions of constant light.....	30
Figure 3.2 Arrhythmic plants show no variation in susceptibility with respect to time of infection under free-running conditions of constant light.....	32
Figure 3.3 Second cycle infections show time-of-day variation in Col-0 but not in <i>CCA1-ox</i>	33
Figure 4 Bacterial counts in Col-0, <i>CCA1-ox</i> and <i>elf3-1</i> after 4 and 48h when infected with (i) <i>P. syringae</i> DC3000 <i>avrB</i> and (ii) <i>P. syringae</i> DC3000 <i>avrRpt2</i> at CT26 and CT42.....	35
Figure 5.1 Callose levels measured at CT26 and CT42 in Col-0 and <i>CCA1-ox</i>	36
Figure 5.2 Fluorescent microscope images showing callose deposits.....	36
Figure 6.1 <i>OX11</i> expression measured at CT26 and CT42 24h after infection with <i>P. syringae</i> DC3000.....	38
Figure 6.2 shows <i>PR1</i> expression measured as bioluminescence at CT26 and CT42.....	39
Figure 6.3 Gene expression of (i) <i>WRKY22</i> and (ii) <i>WRKY29</i>	40
Figure 6.4 Basal levels of <i>SRFR</i> in Col-0 at CT26 and CT42.....	42
Figure 6.5 Expression profiles of <i>CCR2::LUC</i> before and after inoculation with mock (10mM MgCl ₂) or infected (<i>P. syringae</i> DC3000) under free running conditions (LL).....	43

List of Tables:

Table 1 The various strains of *P. syringae* used and the required antibiotics.....24

Table 2 Primers used, annealing conditions and R² values.....27

University of Cape Town

Abbreviations

AOS	Active Oxygen Species
<i>Arabidopsis</i>	<i>Arabidopsis thaliana</i>
AtRBOHD	Arabidopsis Respiratory burst oxidase homologs
BAK1	BRI1 Associated Receptor Kinase 1
CCA1	Circadian Clock Associated 1
CCA1-ox	Circadian Clock Associated 1-overexpresser
CHE	CCA1 Hiking Expedition
Clk	Clock
CLR	CATTERPILLAR
Col-0	<i>Arabidopsis thaliana</i> Columbia-0
CRY	CRYPTOCHROME
CSP	Cold Shock Protein
Cfu	colony forming units
Cyc	Cycle
DBT	Doubletime
<i>Drosophila</i>	<i>Drosophila melanogaster</i>
EE	Evening Element
EFR1	EF-Tu Receptor 1
EF-Tu	Elongation Factor Tu 18, 20
elf18	Elongation Factor Tu
elf26	Elongation Factor Tu
EMT	Early MAMP triggered defence
ETI	Effector Triggered Immunity
flg22	flagellin 22
FLS2	Flagellin Insensitive 2
GLS5	Callose synthase 5

Hpa Emwa1 *Hyaloperonospora arabidopsis Emwa1*

JMJD4 Jumonji Domain containing 4

LHY Late Elongated Hypocotyl

LRR Leucine Rich Repeat

MAMP Microbe Associate Molecular Pattern

ME Morning Element

MTI MAMP Triggered Immunity

NK Natural Killer Cell

OXI1 Oxidative Signal Inducing 1

P. syringae *Pseudomonas syringae DC3000*

PAD4 Phytoalexin Deficient

PCD Programmed Cell Death

Per Period

PGN Peptidoglycan

PMR4 Callose synthase

PR Pathogenesis Related

PR1 Pathogenesis Related 1

PRR Pattern Recognition Receptor

qPCR quantitative PCR

R Resistance proteins

RIN4 RPM1 interacting Protein 4

RLK Receptor Like Kinase

RNAi RNA interference

ROS Radical Oxygen Species

SA Salicylic acid

SAR Systemic Acquired Resistance

SD Sprague Dawley

SID2	Salicylic Induction Deficient 2
SRFR1	Suppressor of RPS4-RLD1
Timeless	Tim
TLR	Toll Like Receptor

University of Cape Town

Chapter 1: Literature Review

For millions of years, plants and pathogens have been engaged in an arms race that led plants to evolve a two-tiered innate immune system [1]. Plant immunity evolved to recognise conserved structural features of pathogens using transmembrane pattern recognition receptors (PRRs) [2,3]. These conserved microbe structures include flagellin and peptidoglycan (PGN) and are collectively termed microbe associated molecular patterns (MAMPs) [4]. Upon recognition of these MAMPs, plants initiate MAMP triggered immunity (MTI) [1]. To overcome MTI, pathogenic bacteria evolved effector proteins some of which suppress MTI responses thus enhancing virulence [1]. These effectors often delivered by a type three-secretion system into the plant cell act to manipulate host defences and suppress MTI [1]. The plant immune system recognises some effectors through resistance protein (R proteins) resulting in effector triggered immunity (ETI), an accelerated and amplified MTI response [1]. R proteins are intracellular and recognise effectors either directly or indirectly with more severe responses such as programmed cell death (PCD) [1]. This continuous conflict between plants and pathogens characterised by bacteria acquiring new effectors to enhance virulence and plants evolving new R proteins to detect them is described as the plant-pathogen arms race [1]. This review investigates the *Arabidopsis thaliana* (hereafter *Arabidopsis*) immune system and compares aspects of plant innate immunity with immune systems of mammals and insects. Current literature purports that immune systems of plants, insects and mammals evolved strikingly similar features by convergent evolution [5,6]. Circadian clock regulation of innate immunity is also common to insects, animals and humans. The immune systems of mice, humans and fruit flies demonstrated some degree of circadian regulation [7,8,9], while recently potential clock control of plant defence was proposed [10]. This review summarises similarities in innate immunity, particularly focussing on circadian regulation of innate immunity in plants, animals and insects. Additionally, I report on recent advances implicating the plant circadian clock in the regulation of immune responses and speculate on additional questions these latest discoveries reveal. Finally, the review concludes with a brief introduction to the questions this study sought to answer, and the experimental approaches taken.

The *Arabidopsis* immune system

The *Arabidopsis-Pseudomonas syringae* DC3000 (*P. syringae* DC3000) interaction has been very useful in developing our understanding of the plant immune system. While the *Arabidopsis*' genome is sequenced thus making it an ideal candidate for molecular studies, *P. syringae* DC3000 is also well

characterised. This two-model system contributed greatly to contemporary understanding of plant immunity.

Plants possess a two-tiered innate immune system consisting of MTI and ETI. Perception of pathogens also results in Systemic Acquired Resistance (SAR) [11]. Using the plant hormone, salicylic acid (SA), plants are able to signal messages away from the site of infection to uninfected cells resulting in activation of defence genes, giving broad spectrum protection to the plant known as SAR [12]. Additionally plants use RNA interference (RNAi) against viruses [13]. However, this review focusses on the MTI and ETI responses in plant-pathogen interactions.

As plants lack mobile defence cells, they rely on innate immunity of individual cells and systemic signals emanating from infected sites [1]. MAMPs are common to most bacteria and because they are essential to bacterial survival and fitness undergo very little change over time [14]. PRRs, which are cell membrane bound receptors on plant cells, recognise these conserved MAMPs leading to MTI [1]. Some bacteria have evolved mechanisms to counter MTI. Bacteria possessing type III secretion systems (TTSS) release effector proteins into the intracellular space of plants, aiming to suppress MTI [1]. In turn, plants evolved R proteins that detect these effectors leading to ETI. Bacteria seek to evade ETI by losing or modifying the effectors in their repertoire, restoring virulence [1]. This zigzag model of plant-pathogen interactions is described as the plant-pathogen arms race [1]. A previously established view of plant innate immunity is that the plant immune system is a two-branched innate system, however more recently experimental evidence summarised in reviews by Tsuda (2010) and Thomma (2011) [15,16], suggest there is considerable overlap between MTI and ETI. The notion that plant innate immunity involving MTI and ETI is a continuum rather than a two-branched system will be discussed in detail later in this review. I will begin by discussing each branch separately.

MAMP triggered immune response (MTI)

The MTI response is initiated by plant recognition of MAMPs [1,6]. Known MAMPs include flagellin (flg22), elongation factor-EF-Tu (elf18 and elf26), chitin and PGN [1,4,17,18,19]. These MAMPs are recognised by PRRs such as FLAGELLIN INSENSITIVE 2 (FLS2), which are Receptor like kinases (RLK) with an extracellular Leucine Rich Repeat (LRR) domain and an intracellular kinase signalling domain [1,4,17,18,19]. Upon recognition of MAMPs, the surface receptor is internalised by endocytosis and activation of defence results [20]. Upon recognition of flg22 by the FLS2 receptor

there is rapid transcriptional induction of 1100 *Arabidopsis* genes [21]. Another MAMP, elongation factor Tu (EF-Tu), is recognised by the PRR EF-Tu Receptor1 (EFR1) [18] and triggers induction of a similar set of genes to flg22, suggesting that although MAMP receptors are different, the downstream defence pathway activated by their recognition is very similar [22,23]. In addition to the MTI response, a weak ETI response is also initiated as some effectors are weakly recognised [1]. Defence responses include strengthening of cell walls (by lignification), callose deposition, and production of hydrolytic enzymes and expression of Pathogenesis Related (PR) proteins such as phytoalexins [24,25,26]. Additionally hormonal changes occur with levels of ethylene, jasmonic acid and salicylic acid all affected [27,28]. These hormones are crucial for plant defence regulation [27,28]. Callose production is a classical marker of MTI after treatment with MAMPs or non-infectious *P. syringae* [29,30]. Perception of fungal and bacterial pathogens elicits callose synthesis by callose synthase GLS5/PMR4 [31,32,33]. Compared to wild type plants, *pmr4* mutant plants showed 20-fold higher counts of *P. syringae hrcC* which lack the type three-secretion system [33], illustrating the importance of the callose response during immune responses. Callose is secreted by the plant in response to infection, slowing down the rate at which bacteria proliferate. This gives the plant time to mount a stronger defence response while slowing down rate of infection.

Early molecular events upon MAMP perception:

FLS2 is a membrane spanning plant LRR receptor essential for recognition of flagellin [34,35]. The intracellular domain of FLS2 interacts with a receptor-like kinase (RLK) called BRI1 associated receptor kinase (BAK1) [34,35]. The interaction between FLS2 and BRI1 is vital for activation of downstream responses after recognition of flagellin as downstream signalling is strongly impaired in *bak1* null mutants [34,35]. Plant recognition of MAMPs results in alkalinisation of the intracellular space within 0.5-2.0 min [36,37,38]. Cytoplasmic levels of Ca^{2+} and other charged ions also increase upon perception of MAMPs and play a critical role in the activation of defence responses [39]. A virulence strategy employed by invading bacteria such as *P. syringae* is to sequester Ca^{2+} ions using bacterial exopolysaccharides to chelate free ions [40]. This is an indication of the importance of Ca^{2+} in defence signalling [36].

In addition to a flux in ions such as Ca^{2+} , production of reactive oxygen species (ROS) which occurs as an oxidative burst is also reported [36]. The ROS signal, generated by membrane bound NADPH oxidases such as *Arabidopsis Respiratory burst oxidase homologs (AtRbohD)*, results in the production of active oxygen species (AOS) such as H_2O_2 , which act as a form of defence as they are

harmful to the invading bacteria [41,42]. AOS also activate Oxidative Signal Inducing 1 (OXI1), a serine/ threonine kinase, which is needed for full activation of the mitogen-activated protein kinase (MAPK) pathway and MTI response [43,44]. Perception of MAMPs such as flg22, chitin and PGN activate the MAPK cascade [45,46,47,48,49]. FLS2 simultaneously activates MPK3/6 via MKK4/5, and MPK4 via MEKK1/2, which act as activators and repressors of defence responses, respectively. While MPK3/6 contributes positively to MTI, MPK4 acts to suppress MTI. These contradictory responses to MAMP perception are mediated by a PP2C phosphatase [36,50] and will be discussed in detail later.

One of the effects of activation of MPK3/6 pathway by MAMP perception is the activation of WRKY22/29 transcription factors that are involved in defence gene expression [45]. The ability of plants to respond to multiple biotic and abiotic signals is achieved through complex molecular pathways that are yet to be fully dissected. However, the family of WRKY transcription factors is a key signalling network involved in numerous processes, most notably in biotic and abiotic stress response [51]. The WRKY family of transcription factors get their name from the highly conserved WRKYGQK peptide sequence and a zinc finger motif present in all members [52]. The large WRKY family shows overlapping functionality among its member proteins for example, WRKY22 is homologous to and shows functional redundancy with WRKY29 [53]. WRKY29 was shown to be a key transcriptional activator of defence genes in *Arabidopsis* innate immunity [45]. WRKY29 induction occurs rapidly, within 30min of MAMP perception whereas defence responses such as *PATHOGENESIS RELATED 1 (PRI)* production occur much later [45]. *PRI* expression is associated with systemic acquired resistance (SAR) and the hypersensitive response (HR), which is mediated by salicylic acid and salicylic acid associated genes such as *SALICYLIC INDUCTION DEFICIENT2 (SID2)* and *PHYTOALEXIN DEFICIENT (PAD4)* [54]. Initially HR and SAR were thought to be characteristic of ETI responses only but recently, MTI was shown to initiate them too [55]. Thus, while the MTI response shares similar downstream molecular pathways as ETI, ETI is initiated by recognition of effectors which are often strain specific and delivered through the type-three secretion system of bacteria into the plant intracellular space [1].

To evade the MTI response, bacteria evolved effectors that suppress MTI responses with the model bacterium *P. syringae* DC3000 secreting more than 30 effectors [1]. These effectors abrogate MTI responses by targeting plant defence molecules for degradation, for example, the bacterial effector AvrRpt2 targets the *Arabidopsis* RPM1 interacting protein (RIN4) for degradation [56]. RIN4 regulates stomata aperture during pathogen attack thus limiting bacterial invasion [57]. Effectors such as AvrPto which is a kinase inhibitor [58] suppresses auto-phosphorylation of PRRs such as FLS2

[59], while AvrPtoB possesses an E3 ligase domain [60] and targets FLS2 for degradation. The presence of effectors abrogates MTI responses but effectors themselves are targets for the plant immune response.

Effector Triggered Immune response (ETI)

Recognition of effectors results in avirulence, a rapid and strong defence response and activation of HR and PCD at a much quicker rate than MTI [1]. Through direct gene-for-gene resistance, the ‘guard’ model and the ‘decoy’ model, plants detect effector molecules resulting in a stronger and more prolonged defence response compared to MTI [1,61]. The ETI response has a biphasic nature to it, with the initial phase triggered by MAMP recognition, which is suppressed by effector molecules [1]. The effector molecules in turn trigger the second phase of plant defence as R proteins recognise them [16,62]. *Arabidopsis* possess approximately 125 R proteins that may recognise effectors secreted by bacteria [63]. Notably, R proteins are race specific and different populations of a single species may have different R proteins [63]. These R proteins, which are intracellular, are important in mediating the effector-triggered response as they detect effectors secreted by bacteria into the cytoplasm [16,36]. This detection can be direct or indirect.

The effectors discussed earlier: AvrPto and AvrPtoB, are capable of suppressing MTI by binding to FLS2 [58,59]. However, resistant *Solanum lycopersicum* (tomato plants) detect the effectors directly using the RLK PTO and the R protein PRF that directly interact to activate ETI [64,65]. The direct interaction of PTO with AvrPto occurs in the presence of PRF, a Nucleotide Binding- Leucine Rich Repeat (NB-LRR) R protein [59]. Interestingly AvrPto only contributes to virulence in the presence of FLS2 suggesting binding of FLS2 and other RLKs are the operative mechanism by which virulence is achieved [59]. In a recent review proposing the decoy model for plant defence, PTO was suggested as a decoy molecule to bacteria expressing AvrPto thus attenuating virulence [66]. PTO is a RLK, similar to FLS2 and competes with FLS2 for AvrPto effectively sequestering the effector before it can bind to FLS2 and other RLKs involved in defence [64]. PTO interacts *in vivo* with PRF and its accumulation is dependent on levels of PRF [64]. This suggests that PTO functions in a dose-dependent manner, outcompeting other RLKs for AvrPto, thus reducing virulence [67]. This is termed the ‘competing decoy’ model [66].

Another way in which plants detect effector activity is via the ‘guard’ model. This model is described in detail by Dangl and Jones (2006) [1] and the key tenets are described briefly. The guard model proposes that bacteria secreting effectors that suppress MTI may alter plant proteins, creating ‘pathogen-induced modified-self’ molecular patterns, which in turn are detected by R proteins leading to ETI. The effector proteins AvrB and AvrRpt2 both interact with RIN4, phosphorylating it and targeting it for degradation, respectively [56,68] effectively modifying the plant molecule. The NB-LRR protein RPM1 detects AvrB activity indirectly, while RPS2 indirectly detects AvrRpt2 activity [68,69]. Thus, RPM1 and RPS2 guard proteins monitor RIN4 and upon modification of RIN4 by effectors, initiate ETI. In the *rpm1rps2* double-null mutant RIN4 is manipulated by bacterial effectors to promote virulence [1]. The effector-triggered immune response also results from direct gene for gene recognition. While ETI is a stronger and more prolonged response, it shares similar molecular pathways to MTI and results in the activation of similar defence responses, hormonal changes and systemic responses. Recent reviews [15,16] excellently show that there is considerable overlap between MTI and ETI. The shared molecular circuitry in *Arabidopsis* suggests that while R proteins are thought to have evolved after PRRs, they evolved to share the same molecular pathways and defence outputs [16].

The two-branched paradigm challenged:

Recent reviews have questioned the basis of the widely held view that the plant immune system possesses two distinct branches of immunity [15,16]. Using data from recent experiments, Thomma *et al.* (2011), illustrated that widely held assumptions about MAMPs, effectors, PRRs and R proteins are all debatable [15]. MAMPs, recognised for their importance in bacterial survival and fitness [70,71], are structural features that are highly conserved across genera, while effectors are specific to single or a few related species [1,15,72]. Apart from their distribution their intrinsic roles are used to differentiate MAMPs and effectors: MAMPs contribute to survival and fitness while effectors specifically contribute to virulence by targeting host defence [15]. Several examples show that it is difficult to separate virulence from fitness and survival [15]. Experiments showed MAMPs such as flagellin and harpin (HrpZ) are essential for virulence [73,74,75,76]. Another MAMP Ax21 released in a cell density dependent manner could be important in quorum sensing and therefore play a role in virulence of bacteria [77]. Perhaps one of the ways MAMPs contribute to survival and fitness is through their role as virulence molecules.

In addition to MAMPs contributing to virulence, they are also able to suppress MTI, a feature previously assigned to effectors alone [15]. Chitin from *Cladosporium fulvum* (*C. fulvum*) activates MTI after recognition by a LysM PRR protein [78]. Using the zigzag model, one could speculate that the fungal pathogen may produce an effector to suppress the resulting MTI response. Indeed the fungus does produce a molecule, Ecp6 that mimics the host Lys M PRR and is a receptor-like protein (RLP) [79]. Ecp6 attenuates the MTI response to chitin by binding to any free chitin before the plant Lys M receptor can, thus sequestering any free chitin [80]. All strains of *C. fulvum* possess Ecp6 with little sequence variation and conserved Ecp6 orthologs are common in the Fungal Kingdom [79]. Its wide distribution across strains of *C. fulvum* and conserved sequence mean it is comparable to a MAMP [15] yet its ability to suppress MTI qualifies it as an effector. Conversely, MAMPs with a narrow distribution have also been described, with only a narrow range of host plants recognising them, a characteristic often attributed to effectors [15].

PRRs and R proteins show distinct characteristics. PRRs are membrane bound, highly conserved and evolutionarily ancient while R proteins are intracellular and evolved more recently [15]. The discoveries of R proteins that display typical characteristics of PRRs further blur the line between MTI and ETI. The R protein Ve1 found in tomato contributes race specific resistance to race 1 strains of vascular fungal pathogens of the *Verticillium* genus [81]. However, the race 1 resistance affects two distinct species of fungus *V. dahliae* and *V. albo-atrum* [15]. This suggests Ve1 mediated resistance is activated by an elicitor common to different fungi, most likely a MAMP, thus Ve1 shows characteristics of a PRR, not an R protein [15]. Ve1 shows additional PRR characteristics such as activating only a weak MTI response [15] and interacting with BAK1 a crucial component of MTI responses [34,35]. A defining characteristic of PRRs is their capacity to recognise broad structural patterns across species but recently close inspection of PRRs revealed some PRRs have a narrow host range [19,22,82,83]. The PRR EFR1 that recognises Ef-Tu is found in *Arabidopsis* but not in *Nicotiana benthamiana* (tobacco) [22] while bacterial cold shock protein (CSP) a MAMP, elicits MTI responses in tobacco but not in tomato, rice and *Arabidopsis* [19]. These results show that only a narrow host range recognises widespread MAMPs. Additionally PRRs that are specific for the same MAMPs recognise different MAMP epitopes implying that some PRRs may have evolved recently thus challenging the assumption that PRRs are evolutionarily ancient [15].

MTI is often characterised as a weak defence response while ETI is characterised as a stronger and more prolonged response. Features thought to be unique to ETI such as HR and SAR have recently been shown to occur in MTI [55,84]. In addition to strong MTI responses, weak ETI responses also occur for example RPS4 resistance to AvrRps4 in *Arabidopsis* [85].

Thomma *et al.* (2011) present an excellent review citing numerous examples all pointing towards the conclusion that a continuum exists between MTI and ETI responses. They conclude that the robustness of MTI or ETI depends on the specific interaction; different molecules activate defence pathways of varying strength, depending on the trigger, the receptor and possibly environmental conditions [15]. Interestingly, they also suggest that plant innate immunity evolved to regulate the strength of defence depending on the type of microbial attack. Ausubel (2005) [6] introduced a similar idea in a review where he suggested that the two-tiered response might have evolved as a mechanism to distinguish between non-pathogenic and pathogenic bacteria. As immune responses are energetically expensive, this 'two hit' approach to defence would allow a moderate defence response to less harmful bacteria and full response would only manifest in the event of infection by pathogenic bacteria. The two-branched model and zigzag model while distinctly separating the defence responses, acknowledged that an overlap might exist. Recent findings support the view that a continuum exists between the MTI and ETI responses and as our understanding of plant immunity grows these two previously separate branches may merge even more.

Negative regulation of plant defence

Plants are able to distinguish between pathogenic and non-pathogenic bacteria suggesting careful control of defences. This is important as a full immune response is energetically expensive and comes at great cost to the plant often resulting in cell death. Understanding negative regulation of plant immunity will help to explain how plants co-ordinate defence responses to deliver a proportionate response.

Genetic studies revealed a number of negative regulators of various aspects of immunity: SNI1 represses systemic acquired immunity, LSD1, and ACD11 prevent programmed cell death [86]; and SRFR1 regulates levels of NB-LRR R proteins preventing them from over-accumulating [87], thus avoiding autoimmune responses. Other negative regulators include a number of *WRKY* genes, KAPP that negatively regulates FLS2 and PUB22, 23 and 24. The PUBs negatively regulate MTI possibly by binding to PRRs and degrading positive regulators of MTI [88]. Genetic studies revealed that the plant innate immune system is under negative regulation prior to MAMP perception [89]. However, negative regulation not only occurs in the absence of infection but persists even after activation of defence.

Identification of MAMPs leads to activation of both MPK3/6 and MPK4, which positively and negatively regulate MTI, respectively. PP2C phosphatase AP2C1 regulates both pathways possibly to ensure a sensitive and controlled defence response upon MAMP perception [90]. This may allow plants to distinguish pathogens from commensal microbes [6]. Recently, a comprehensive study, which divided plant defence responses into sectors based on positively co-ordinated pathways, identified four sectors: The Early MAMP Triggered (EMT) genes, the Salicylic Acid-associated (SA) genes, the Jasmonic Acid genes (JA) and Ethylene associated genes [54]. Early MAMP triggered proteins include MPK3/6, WRKY29, PMR4 (callose synthase) and AtRbohD NADPH oxidase. The proteins that make up the SA sector include PAD4, SID1, PR1 and NPR1 which are involved in SA signalling thus activating SAR and HR [54]. While both sectors positively contribute to plant defence, this study reported that the EMT sector negatively regulates the SA sector and that this relationship is antagonistic [54]. For example, PMR4, which is part of EMT, negatively regulates SID1 and *pmr4* null mutants show greater SA mediated activity [32]. Collectively these results suggest that PMR4 negatively affects SA mediated responses and *vice versa* [32]. MPK3/6, Jasmonic Acid and Ethylene all showed negative relationships with the SA sector [54].




Figure 1. The ‘two-hit’ model of plant defence: An illustration of the EMT and SA sectors which show an antagonist relationship. Phase 1 is marked by the recognition of MAMPs such as flagellin and effectors by PRRs and R proteins. This activates the Early MAMP Triggered (EMT) sector. A strong EMT response (shown by a larger circle) clears infection while simultaneously suppressing the SA sector (thicker arrow). Importantly, EMT responses are not harmful to the plant. In the event that the bacteria possess effector molecules capable of suppressing EMT, phase 2 is initiated. Suppression of EMT by bacterial effectors inadvertently activates the SA sector as the antagonistic repression of SA by EMT weakens (with a weak EMT shown by thinner arrow). A strong SA sector in phase 2 (larger circle) also represses EMT in phase 2 and initiates more severe defence responses such as programmed cell death. This model for infection shows how plant defence responses are hierarchical, reserving the most severe response for instances when EMT genes cannot clear infection. Additionally, this model shows how repression of EMT by bacteria results in even stronger responses.

These results indicate that activation of one defence pathway results in repression of another, even though collectively they aid in defence against pathogens [54]. This apparently paradoxical phenomenon points to the need for a nuanced look at plant immunity. Mounting evidence suggests that plants employ a carefully moderated defence response to bacteria [15,16,54]. This restrained approach may allow plants to distinguish between non-pathogenic and pathogenic bacteria thus reserving a full defence response only when necessary. Additionally by dividing defence responses into mutually antagonistic sectors, plants evolved an energetically favourable method of clearing infection. The *pmr4* null mutants that are unable to initiate a callose defence response display a compensatory effect by showing greater SA mediated responses [32].

This finding is important as it points to a hierarchical defence response of increasing severity. Early defence responses such as callose production, the production of antimicrobial peptides and production of AOS are not harmful to the plant compared to defence responses activated by SA, which eventually result in programmed cell death. The antagonistic relationship between the EMT sector and the SA sector is important in plant defence responses (Fig 1). When this antagonistic relationship is perturbed as in the case of *pmr4* null mutants lacking callose synthase, a weak EMT response ensues and so the EMT sector weakly represses the SA sector, which results in a stronger SA sector response. In turn, a strong SA sector response represses the EMT sector.

With these observations, a new approach to understanding events in immune response becomes apparent. PRRs and intracellular R proteins recognise pathogenic bacteria, resulting in a strong EMT response. As a result, a strong EMT response has an antagonistic effect on the SA sector thus repressing it. This is phase 1 of infection (Fig 1). In the event that the bacteria possess effectors or MAMPs capable of suppressing EMT, phase 2 is initiated. Suppression of EMT by bacterial effectors inadvertently activates the SA sector as a weak EMT response is only weakly antagonistic to the SA sector (Fig 1). Additionally, a stronger SA sector has an antagonistic effect on the EMT sector, further repressing it. Thus, by overcoming early defence responses virulent bacteria switch on the SA sector, activating stronger defence responses and SAR. The ultimate effect of the SA sector is programmed cell death, which is costly to the plant but necessary to isolate the pathogen at the site of infection. This model offers a new explanation of events in defence responses to pathogenic and non-pathogenic bacteria and could be described as the ‘two-hit’ approach to plant defence, a term coined by Ausubel (2005) [6]. Furthermore, this model explains how plants may keep non-pathogenic and commensal microbes at bay, without having to sacrifice large resources or initiate strong defence responses that

are detrimental to the plant. Commensal bacteria may trigger EMT as they possess MAMPs but lack effectors or MAMPs that can suppress EMT responses and so a full-blown defence response is not required. Since the EMT sector is not repressed, it has an antagonistic effect on the SA sector preventing extreme defence responses to harmless bacteria. The ‘two-hit’ model proposed here is largely similar to the zigzag model. However, the zigzag model proposes that the perception of effectors triggers a stronger response called ETI but it does not explain how that may occur. The ‘two hit’ model shows that suppression of EMT by effectors and possibly MAMPs, is what triggers stronger defence responses, as weakening of EMT has an antagonistic effect on the SA sector (Fig 1). I base the ‘two-hit’ model on ideas reported by Thomma *et al.* (2011) and experiments by Sato *et al.* (2010) [15,54]. As some MAMPs can suppress EMT genes [40], the ‘two-hit’ model also explains why MAMPs were reported to trigger HR and PCD [55].

Comparing innate immunity in plants, animals, insects and invertebrates:

Plants, animals, insects and invertebrates such as *C. elegans* possess innate immune systems. Plants and animals share a number of common features in their immune systems for example both recognise MAMPs such as flagellin and harpin using membrane bound PRRs [6]. This observation initiated a debate questioning whether these similarities were due to divergent evolution or convergent evolution. A number of scientists have commented on whether the shared features of innate immunity found across kingdoms are the result of divergent evolution from a common ancestor or the result of convergent evolution [91,92]. A number of reviews as well as mounting experimental evidence indicate that common features of innate immunity arose independently in each kingdom and are therefore best explained by convergent evolution [6,14,37]. Owing to a limited set of molecules and being subject to the same environmental constraints perhaps led plants and animals to independently evolve analogous approaches in dealing with microbe attack [6].

The most obvious parallel between plant and animal innate immunity is the presence of defined receptors for detecting MAMPs. Both plants and animals share similar PRRs which recognise MAMPs such as flagellin and both animal and plant PRRs activate defence signalling via a MAPK signalling pathway [6]. In plants, the LRR protein FLS2 recognises flagellin. The analogous PRR in animals is the toll like receptor 5 (TLR5) [6]. These PRRs share a number of similarities: they are membrane bound receptors with Leucine rich regions and upon recognising flagellin initiate defence responses [6]. However, closer inspection revealed that these PRRs recognise different epitopes of flagellin [93]. Additionally, the LRR domains are very divergent and plant FLS2 lacks an intracellular

TIR domain and other downstream components present in TLR5 [6]. Additionally, FLS2 has an intracellular kinase domain whereas TLRs recruit IRAK kinase for downstream signalling [6]. Flagellin perception in plants results in activation of defence via WRKY transcription factors which are absent in animals and insects [6]. Animals and insects evolved the transcription factors NF- κ B and Dif respectively to activate transcription of antimicrobial peptides [6].

Both plants and animals produce an AOS signal upon MAMP perception and is mediated by gp91^{phox}-like NADPH oxidase (AtRbohD in plants) common to both [94,95,96]. This observation could support the view that shared components of plant and animal innate immunity suggest a common ancestor. However, gp91^{phox}-like NADPH oxidase is also involved in a number of other biological processes in both plants and animals and therefore could have been recruited to pathogen defence independently [6]. Lastly, the best-characterised plant defence molecules are the R proteins (NBS-LRR) which possess a tripartite structure similar to animal CATERPILLAR (CLR) also called Nod proteins [97,98]. While both classes of defence molecules initiate programmed cell death as a defence mechanism, there is no evidence suggesting common signalling components [6]. Collectively, these examples show innate immune systems share similarities but these similarities are likely due to plants and animals converging on similar solutions to similar problems.

The similarities in the innate immune systems of animals, insects and plants suggests a common 'logic' to the overall structure of innate immunity in these kingdoms [6]. Studies on plant immunity revealed novel insights into animal innate immunity and *vice versa* and continue to do so. Advances in mammalian and insect immunity showed the circadian clock contributes to regulation of innate immunity. The plant circadian clock contributes to general fitness and recent experimental evidence linked a component of the plant circadian clock to plant innate immunity. Understanding similarities among the innate immune systems of animals, plants and insects is essential in getting a clearer picture of the inner workings on innate immunity.

The Circadian Clock and Immunity

The circadian clock regulates a number of biochemical and physiological processes in plants, fungi, animals and insects. The Earth rotates on a fixed axis, which results in a 24h day with light and dark cycles of varying length. Over evolutionary time, it is widely thought that organisms independently evolved circadian clocks to regulate various metabolic and physiological processes [99]. Circadian

clocks confer an evolutionary advantage as they enable organisms to anticipate regular daily events, thus priming various physiological processes to coincide at the most favourable time [99,100,101]. Circadian clocks are characterised by three features: they are regulated by transcriptional-translational feedback loops, can be reset by light and temperature input and show temperature compensation and continue to run under constant conditions [100,102,103,104]. The ability of circadian clocks to reset is essential as it allows organisms to synchronise their internal processes with the external environment, adapt to changing day length and anticipate seasonal change [99]. Temperature compensation is also an important characteristic of the circadian clock. Usually for a 10°C, increase in temperature there is an associated increase in biochemical rate of activity. If the circadian clock was not temperature compensated, it would run faster at higher temperature and slower at lower temperature [99]. However, the 24h period of the clock is largely unaffected by temperature and this is reflected by circadian clocks retaining near 24h rhythms at varying temperatures, which is important in organisms that cannot regulate their own temperature [102,103]. Circadian clocks are essential for a number of processes and have been demonstrated to play a role in immune responses in flies, mice, humans and plants.

The Circadian Clock of *Drosophila melanogaster* and its role in immunity

The circadian clock of *Drosophila melanogaster* (*Drosophila*) the fruit fly maintains a 24h period by regulating expression of *Clock* (*Clk*), *Cycle* (*Cyc*), *Period* (*Per*) and *Timeless* (*Tim*) [99]. Using at least two interconnected transcriptional-translational feedback loops, *Drosophila* is able to retain rhythmic expression of a number of output pathways [105]. PER and TIM associate together, interacting with the CLK-CYC complex and block it from transcribing their expression forming a negative feedback loop [106,107]. This leads to down regulation of *Per* and *Tim* transcription and activation of *Clk* transcription [107,108,109,110]. Transcription of *Per* and *Tim* begins around mid-day with RNA levels peaking in the early evening and protein levels peaking at midnight [110]. While PER is phosphorylated by a kinase DOUBLETIME (DBT), which targets it for degradation, TIM is degraded in a CRYPTOCHROME-dependent manner by light [111,112]. At night, TIM binds to PER and this prevents DBT from phosphorylating PER, thus PER avoids degradation [111,113,114]. When TIM and PER associate they are able to localise to the nucleus [114,115,116]. Nuclear localised TIM-PER undergo phosphorylation events, which eventually mark the complex for degradation [110,117]. After degradation of TIM-PER repression of CLK-CYC ceases resulting in activation of *Per* and *Tim* transcription (the positive feedback loop) and repression of *Clk* transcription [118]. These interconnected feedback loops form the core rhythms of the circadian clock in *Drosophila*.

Lee and Edery (2008) showed circadian regulation of *Drosophila* immune response to *Pseudomonas aeruginosa* and *Staphylococcus aureus*. To investigate clock control of immune responses in *Drosophila*, infections were performed at various time points under light-dark cycles and in free running conditions of constant dark [119]. Circadian rhythms persist under constant light or dark and constant temperatures, known as free-running conditions. Flies showed greater susceptibility to the bacterial pathogens after daytime infections compared to infections at night. The time-of-day variation to infection was also observed under constant dark conditions suggesting circadian clock regulation of defence responses [119]. To test this, experiments using clock mutant flies: *Per⁰¹*, *Clk^{Jrk}*, *Cyc⁰¹* and *Tim⁰¹*, showed that loss of clock components resulted in loss of time-of-day variation in susceptibility to infection by *P. aeruginosa* [119]. Interestingly, the clock mutants behaved differently to infection, some showing enhanced defence responses while others were more susceptible to infection. This suggests loss of different components of the clock affected immune responses uniquely [119]. The *Per⁰¹* mutant showed greater susceptibility to infection while *Clk^{Jrk}* was less susceptible than the wild type [119]. *Per⁰¹* mutants also showed greater susceptibility than wild type flies when infected with *Streptococcus pneumoniae* and *Listeria monocytogenes* [7].

In contrast the clock mutant *Clk^{Jrk}* showed lower bacterial counts than *Per⁰¹* with bacterial levels comparable to wild type flies that were infected at night [119]. Collectively these observations implicate clock control of defence genes leading to the *Drosophila* immune system being better primed for defence at night than during the day. Microarray data revealed that basal levels of a key regulator of defence called *imd* showed time-of-day variation in expression with peak expression at night [120] confirmed by quantitative PCR [119]. Additionally, both PGRP-SA a PRR essential for detecting peptidoglycan and *drc* a gene encoding antibacterial peptides accumulate to greater levels at night than during the day [119]. Interestingly the *Per⁰¹* mutant accumulated greater amounts of *drc* than *Clk^{Jrk}* despite *Per⁰¹* showing greater susceptibility [119]. This contradictory observation indicates an instance where incorrect or over-expressed immune genes can have deleterious effects. Overexpression of defence genes perhaps abrogates other defence responses while also being physiologically expensive [119,121,122,123]. This in turn leads to diminished overall fitness and greater susceptibility to infection. Lee and Edery (2008) illustrated the role of circadian rhythms and clock genes in immunity. Their work suggests PER may play a protective role in innate immunity by regulating expression of defence genes.

Circadian regulation of immune responses in mammals

Mammalian clock components share remarkable sequence similarity with those of *Drosophila* and similarly the clock functions by transcriptional-translational feedback loops. The positive arm of the clock involves CLOCK:BMAL1 associating to regulate expression of *Per 1,2* and *3*, *Cry 1* and *2* and *Bmal1* while PER 1 and 2 and CRY1 and 2 constitute the negative arm of the clock [124]. CRY1 and CRY2 interact with PER1 and PER2, enabling nuclear translocation where they repress CLOCK activity [124]. Like in *Drosophila* PERs 1, 2 and 3 are phosphorylated repeatedly, eventually marking them for degradation [99].

Natural killer (NK) cells are an integral component of mammalian innate immunity [70]. NK cell activity displayed circadian rhythms when investigated under free running constant dark conditions [8]. NK cells produce cytolytic factors and cytokines in response to infection and these molecules play a vital role in NK activity [8]. Expression of the cytolytic factors: granzyme B and perforin, as well as cytokines: IFN- γ and TNF- α , all showed rhythmic expression in NK cells of Sprague Dawley (SD) rats, peaking in expression at night [8]. Additionally, RNA- mediated knockdown of *Per2* resulted in reduced levels of granzyme B and perforin levels [8].

NK cells are essential for detection of infected and malignant cells [70]. NK cells destroy infected cells by calcium-dependent release of cytolytic granules and activation of death receptors in target cells by TNF-related ligands [8]. When NK cell activity in SD rats was investigated for time-of-day variation, maximum activity was reported at night [125]. The collective results show NK cell activity in SD rats is under clock control and peaks at night. SD rats are nocturnal and are most active at night. It is possible that NK cell activity increases at night when exposure to pathogens and other environmental insults would be greatest [8]. That loss of *Per2* resulted in reduced levels of granzyme B and perforin suggests *Per2* positively contributes to production of cytokines and cytolytic factors.

A well-characterised output of the circadian clock of animals is the sleep-wake cycle [126]. An excellent review documents how infections in animals and humans leads to altered sleep patterns, an extreme example is sleeping sickness (*Trypanosomiasis*) caused by a tick-borne pathogen [9,126]. The link between altered sleep patterns after infection by pathogens suggests that the human immune system may also show circadian regulation. Studies revealed that levels of lymphocytes and monocytes reach maximal levels around midnight while NK cells reach peak levels in the afternoon,

decreasing in number at midnight [127,128]. These daily fluctuations are under circadian control, showing a link between sleep and disease response [9]. The effect of sleep on immune responses is beyond the scope of this review but an excellent treatise discussing it can be found here [9]. In summary, the circadian clock regulates immune function of fruit flies, rats and humans and probably other animals.

The Plant Circadian Clock and its role in Immunity

Plants are sessile organisms and unlike animals lack mobile defence cells, rather evolving to have innate immunity in each cell [1]. Similarly, each cell possesses the molecular circuitry that makes up the circadian clock [100]. The synchronised expression of circadian clock components in each cell gives the plant a collective rhythm [129]. In plants, the circadian clock is essential for a number of physiological and biochemical processes including photosynthesis, flowering and seed germination. As in animals and insects, the plant circadian clock is temperature compensated, can be reset by light input and possesses multiple transcription-translational feedback loops [100]. Plant circadian rhythms have a 24h period and persist under free-running conditions [100]. While the circadian clocks of animals, plants and insects share a number of similarities, the actual components of the plant clock are not conserved across the kingdoms and the presence of similar clock circuitry may be a result of convergent evolution [130,131].

While the basics of clock regulation in plants are now known, understanding components of the plant circadian clock is still the subject of active research. Additionally, identifying output pathways regulated by the circadian clock is also researched extensively. Using the model plant, *Arabidopsis*, molecular biologists have managed to dissect the molecular pathways that drive the positive and negative loops of the plant circadian clock [131]. Studying plant circadian rhythms and its clock components are useful not just for the advancement of understanding plant circadian rhythms but also for advancing our understanding of the circadian clocks of animals and insects. Plants, animals and insects share a common architecture among their clocks. Therefore, findings in one system may have implications in another.

The interrelated nature of clock components can be seen in the following example [130]. A histone demethylase jumonji domain containing 5 (JMJD5) which is found in plants and humans [130]. In both *Arabidopsis* and humans, knock down mutants lacking *JMJD5* have a fast-running circadian oscillation phenotype compared to wild type. Intriguingly, the *Arabidopsis* and human JMJD5

ortholog retain enough similarity to rescue the mutant phenotype in reciprocal systems [130]. The human cell line U2OS-B6 has a fast-running oscillation and as it lacks human *JMJD5* [130]. To test if *Arabidopsis AtJMJD5* could rescue the human phenotype, an *AtJMJD5* + U2OS-B6 cell line was created that stably expressed the *Arabidopsis* gene [130]. This cell line was transfected with human short interfering (si) RNA *JMJD5* and the period length measured showing a significantly longer period [130]. However, this was only a partial rescue as further experiments showed a modest decrease in period length in the presence of human siRNA coding *JMJD5* compared to scrambled siRNA [130]. Jones *et al.* (2010) also tested whether the human *JMJD5* ortholog was functional *in planta*, showing that under the control of the *Arabidopsis* promoter human *JMJD5* indeed can restore a longer period.

Like animals and insects, the plant circadian clock has multiple, interlocking feedback loops. The core loop that was initially identified comprises two Myb transcription factors- CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY)- and a pseudo-response regulator called TIMING OF CAB EXPRESSION 1 (TOC1) [131,132]. CCA1 and LHY act together in another loop as positive regulators of PRR7 and PRR9 and this pair in turn, indirectly repress *CCA1* and *LHY* expression [131]. In the core loop, CCA1 and LHY bind together to repress *TOC1* expression and by doing so form the negative arm of the circadian clock [132]. On the other hand, TOC1 positively regulates *CCA1* and *LHY* [132]. Interestingly TOC1 lacks DNA binding motifs and does not directly regulate CCA1 and LHY. Recently, CCA1 Hiking Expedition (CHE) a TCP (TB1, CYC, PCFs) Myb-like transcription factor that binds TOC1 was identified and could aid in understanding the positive limb of the circadian clock [133]. Additionally CHE binds to the *CCA1* promoter repressing *CCA1* expression. Therefore, TOC1 may bind to free CHE, sequestering this repressor of *CCA1* enabling transcription of *CCA1* [131]. This aspect of the core loop is still the subject of current research as our understanding of the positive arm of the plant circadian clock is still incomplete [131]. CCA1 and LHY form homo and heterodimers driving transcriptional processes in various output pathways [134,135] such as expression of the evening phased *AtGRP7* also called *COLD AND CIRCADIAN REGULATED 2 (CCR2)* [136].

As plants are sessile, the circadian clock is crucial for fitness as it enables plants to anticipate regular biotic and abiotic changes, thus priming appropriate metabolic processes to occur at optimal times [100]. A number of plant physiological and biochemical processes are optimal when their internal clock matches the periodicity of the external environment [100]. A short period mutant *toc1-2* when grown under 28h day conditions (14h light and 14h dark) showed reduced fitness with stunted growth and decreased chlorophyll levels, while a long period mutant *ztl-27* grown under 20h days (10h light

and 10h dark) also showed similarly stunted features and overall reduced fitness [101]. These observations indicate the importance of matching the internal clock with the external environment.

Microarray experiments suggest that approximately one-third of expressed *Arabidopsis* genes are under circadian regulation [137]. Analysis of promoter regions of expressed genes also suggests 36% of *Arabidopsis* genes are under clock control [138]. These data suggest the *Arabidopsis* circadian clock regulates a wide range of physiological processes that occur at various times during the light-dark cycle. The genes encoding clock proteins are classified as morning and evening-phased genes depending on when their peak expression occurs. These clock proteins can bind to the promoter region of downstream genes, activating or repressing gene expression at various times of day [100]. The dawn-phased proteins, CCA1 and LHY, bind the Evening Element (EE) found in the *TOC1* gene, resulting in repression of *TOC1* [139,140,141,142]. Dusk-phased genes display enrichment of the EE in their promoters, while dawn-phased genes show overrepresentation of the morning element (ME) [139,141,142]. In addition to ME and EE, PBX regulates expression of night-phased genes and GATA (G-box), is important for expression of late-morning and afternoon-phased genes [139,141,142,143]. Through these various upstream promoter elements, plants may regulate multiple physiological processes to occur at appropriate times during the day.

Plant circadian regulation through complex feedback loops integrates multiple signals thus switching on output pathways at the most advantageous time of day [100]. In addition to generating rhythmic expression of biological pathways, the circadian clock also regulates a number of responses through a mechanism called gating [144]. Gating of gene expression by the circadian clock is a phenomenon where application of a particular stimulus elicits a response only at a specific time in the 24h day [100]. Therefore, a stimulus of equal intensity applied at different times will elicit a response of varying amplitude depending on when the stimulus is applied. An example of circadian gating is shade avoidance in plants [144]. Plants are able to perceive shade in the form of a change in the equilibrium between far-red and red light, through phytochromes [145] with a low ratio of red light to far-red light triggering shade avoidance in plants. Investigating shade avoidance mechanisms in *Arabidopsis* seedlings revealed that 4-day-old seedlings switched on shade avoidance mechanisms such as hypocotyl elongation, in a time-of-day dependent manner [145]. In controlled experiments, plants were transferred from 12h light-dark cycle to constant low red to far-red light mimicking shading but shade avoidance mechanisms were only activated at subjective dusk [144]. Through a gating mechanism, shade avoidance pathways were only activated at dusk even though the stimulus necessary to activate shade avoidance was present throughout the day [144]. Thus, when the circadian clock gates gene expression, the stimulus alone is insufficient to elicit a full response: rather the

stimulus applied at the correct time activates a full response. Gating by the circadian clock perhaps enables plants to regulate physiological processes to occur at the most optimal time of day, ensuring that costly metabolic processes are initiated only at the most opportune moment.

Microarray studies implicate circadian clock regulation of plant responses to biotic stress, suggesting some circadian clock control of plant defence responses to pathogens [137], this idea was clearly articulated by Roden and Ingle (2009) [10]. A number of defence genes have shown circadian regulation and recently a molecular link between the circadian clock and immune responses was established [146].

Seasonal fluctuation in disease in plants is a well-understood phenomenon. Success of plant pathogens in infection is dependent on both host susceptibility and favourable environmental conditions. An example is the release of spores by *Phytophthora ramorum* and *Fusarium circinatum*, which occurs in cool and humid conditions [90,147]. The temporal variation in biotic stress also extends to bacteria on plant leaves showing diurnal variation in abundance [148]. These observations indicate that favourable conditions for infection vary and possibly plants have evolved mechanisms to anticipate them. Indeed a number of plant defence genes display rhythmic expression under the control of the circadian clock [149,150,151,152]. Additionally the defence gene *DEAI* in tomato shows rhythmic expression in long days but is constitutively expressed in short days, suggesting seasonal influence on defence gene expression [10,149]. While these defence genes display circadian rhythms, it would be interesting to investigate whether their induction is regulated by circadian clock gating.

A well-characterised output under clock control is the opening and closing of stomata [153], pores found on the abaxial side of plant leaves. These pores are important for gaseous exchange but are also exploited by bacteria as a portal of entry into the plant leaf. Novel work has revealed that stomata may present a barrier to pathogen entry [154]. Experiments showed that stomata close upon perception of MAMPs [154] while certain bacteria secrete a molecule called coronatine that can re-open closed stomata [154]. Further experiments showed that RIN4 (a target of a number of bacterial virulence effectors) regulates stomata opening and closing during pathogen attack [57]. Regulation of stomata is under the circadian clock and it would be interesting to investigate whether stomata sensitivity to MAMPs and coronatine varies at different times of day as plants attempt to balance the requirements of photosynthesis and pathogen resistance. The clock controlled AtGRP7 sometimes called CCR2

also plays a role in stomatal regulation [155] and is under the control of the circadian clock showing peak expression in the evening [136]. CCR2 is also a target of bacteria with the virulence effector HopU1 affecting the ability of CCR2's to bind to RNA [156]. As CCR2 has been shown to regulate a number of biotic and abiotic stress responses [136], this result suggests that bacteria have evolved mechanisms to target circadian clock output pathways that contribute to defence responses.

Recently, seminal work by Wang and colleagues (2011) [146] established a molecular link between plant defence and a component of the circadian clock called CCA1. Using a systems approach, they identified novel R genes and MAMP genes as they sought to understand how these sets of genes are activated in response to biotic stress [146]. Inadvertently they noticed enrichment of CCA1 and EE binding motifs in the promoter regions of these genes. The R gene *RPP4* involved in plant immune response displayed a circadian rhythm and possessed CCA1 binding elements in its promoter. Wang *et al.* 2011 showed a number of defence genes were potentially under clock control. Using the null mutants' *ztl4* and *cca1* they showed that plant defence responses are compromised in clock mutants. Additionally the over-expressing *CCA1* showed enhanced defence responses to infection. Furthermore, they showed that under alternating light-dark conditions over expression of *CCA1* contributed positively to defence against *Hyaloperonospora arabidopsis* *Emwa1* (*Hpa Emwa1*). Since a number of defence genes possessed a CCA1 binding domain, Wang and colleagues investigated whether *CCA1* expression increased upon infection [146]. Promoter activity of *CCA1* was measured by using the reporter gene luciferase fused to the *CCA1* promoter. Interestingly, *CCA1* expression showed rapid induction and became arrhythmic after infection with *Hpa* [146]. This suggests a *CCA1* mediated defence pathway but further work is required to establish this molecular pathway [157].

Wang and colleagues showed that plants anticipated dawn infection, which corresponded with the release of spores by *Hpa* suggesting that the circadian clock regulated defence responses to occur when perceived threat was greatest. Conversely, greater pathogen success was reported after infection at dusk. This time-of-day variation was absent in the *cca1* mutant [146]. These results show time-of-day variation in the wild type and not in the mutant and they mirror studies performed on *Drosophila* discussed earlier [119]. However, these experiments should be carried out under free-running conditions to investigate whether there is circadian clock regulation [10,119,146,157]. Nonetheless, their novel results provided a link between plant defence responses and the circadian clock showing that absence of a functional circadian clock compromises plant defence.

A systems approach as used by Wang *et al.* (2011) [146] is most likely to yield mechanistic insights into the *Arabidopsis* circadian clock network and its plant innate immune network. Additionally network studies similar to those employed by Sato *et al.* (2010) [54] will also be useful in understanding the relationship between the plant circadian clock and its immune responses.

Concluding remarks

Innate immunity and circadian clock regulation arose independently in insects, plants and animals. These complex systems share similarities across species that perhaps arose because of convergent evolution. The two systems of innate immunity and circadian clock regulation are both the subject of constant research and more recently, the link between the two is gaining greater attention. As our understanding of plant innate immunity grows, a clearer picture emerges which shows that plants attack pathogens with a carefully coordinated defence response. While previous research focussed on innate immunity simply as an on and off system, recent advances indicate that the plant innate immune system is more subtle than that [54]. Implicating the circadian clock in plant defence responses lends weight to the view that plant immunity is carefully coordinated and that plant defence responses may follow a diurnal rhythm showing peak defence when pathogens are most anticipated. The plant defence response is an energetically costly process that affects plant physiology and metabolism irreversibly if fully activated. Therefore, plants perhaps have evolved mechanisms to strike a balance between ensuring protection against pathogens while avoiding the detrimental effects of full-blown immunity. One of the mechanisms could be regulation by the circadian clock and this needs further research.

Research question

The hypothesis that was tested was that defence responses in *Arabidopsis* are regulated by the circadian clock. This work tested the hypothesis by investigating the outcomes of infection by virulent *P. syringae* DC3000 under constant light and temperature conditions. Infections were carried out every four hours over two circadian cycles so that if differences in susceptibility were detected, they could be attributed to endogenous factors. Additionally the arrhythmic clock mutant *elf3-1* and the *CCA1*-overexpressing line were infected with *P. syringae* DC3000. These plant lines are arrhythmic under constant light conditions, thus enabling one to distinguish whether the circadian clock regulates time-of-day variation in response to infection. In addition to using bacterial counts as a measure of plant defences, callose production in response to infection of wild type *Arabidopsis* Col-0 and *CCA1*-

ox with *P. syringae* DC3000 *hrpA* was also measured at subjective day and night. This was used as a measure of the strength of the MTI response. To investigate circadian regulation of specific R protein triggered defence responses wild type, *elf3-1* and *CCA1-ox* were infected with avirulent strains of *P. syringae* DC3000 *avrB* and *P. syringae* DC3000 *avrRpt2*. Promoter activities of *OXII* and *PR1* after infection with *P. syringae* DC3000 at different times of day were investigated using luciferase reporter constructs and transcript levels of *WRKY22* and *WRKY29* were measured in an attempt to understand the basis for differential responses depending on time-of-day of challenge.

University of Cape Town

Chapter 2: Materials and Methods

2.1 Plant Materials

All plant lines used for pathogen assays were in the Columbia background (Col-0). Wild type *Arabidopsis thaliana* were from the Arabidopsis Biological Resource Centre and Seed Stock Centre (<http://www.biosci.ohio-state.edu/pcmb/Facilities/abrc/abrchome.htm>) while the *CCA1-ox* seeds were a kind gift from Dr Alex Webb at Plant Sciences, Cambridge, England. Seeds with *elf3-1 CAB::LUC* were obtained from Dr Frank Harmon, University of California, Berkeley. This *elf3-1* line was used as it was originally made by introgressing the *CAB2::LUC* reporter from wild type C24 into Columbia. This was then backcrossed seven times and while there may be a small amount of C24 around the reporter, the majority of the genome is derived from Columbia (personal communication, Dr. Bryan Thines, UC Berkeley). To measure the promoter activity of *PR1 Arabidopsis Col-0 PR1::LUC* constructs from [158] were used. Promoter activity of *OXII* was measured using *OXII::LUC* in the Col-0 background. *Arabidopsis* in the Wassilewskija (WS) background was used for *CCR2::LUC* promoter activity measurements and were kindly donated by Dr. Seth Davis (Max Planck University Institute for Plant Breeding Research, Cologne, Germany). *OXII::LUC* plants were a contribution from Dr Robert Ingle. The *elf3-1* null mutant and *CCA1-ox* lack a functional clock meaning they are arrhythmic under free-running conditions of constant light and temperature. Many other clock mutants retain some degree of rhythmicity albeit with altered period. Using *CCA1-ox* and *elf3-1* meant that the role of the clock in regulating defence responses could be investigated, rather than a particular clock component.

2.2 Plant growth conditions

All seeds were sterilised for 5 min in 2% (v/v) sodium hypochlorite and 0.025% (v/v) Tween 20 solution. Seeds were washed three times in sterile distilled water and placed on Murashige and Skoog (MS; Highveld biological PTY, Ltd., Lyndhurst, South Africa) medium, pH 5.8 solidified with 0.8 % (w/v) agar. All seeds were stratified at 4°C in the dark for two to three days after which, they were placed in a growth room with long day photoperiod (16h light, 8h dark) with cool white light (60-100µmol m⁻²s⁻¹) at 22°C and 55% relative humidity, except for *CCR2::LUC* plants which were grown under 12h light dark cycles. The seedlings grew on MS agar for seven days and thereafter were transferred to Imidocloprid-treated ('Gaucho', Bayer, Paarl, South Africa) soil which was composed of peat (Jiffy Products, International AS, Norway) and vermiculite in a 1:1 (v/v) ratio. Plants were grown for three to four weeks in plant growth rooms using earlier mentioned conditions. Plants were

moved, within the light period, from long days (LD), to constant light (LL) at least 24 hours prior to pathogen treatment. An explanation of circadian time is shown in Fig 2.




Figure 2. An illustration explaining circadian time and free-running conditions. Plants were grown for 4 weeks in alternating light-dark cycles of 16h and 8h respectively. After that, they were transferred to constant light and temperature conditions. Once in constant light, time is measured as circadian time (CT). Plants were infected at CT26 and then every four hours thereafter, i.e. CT30, CT34 and so on up to CT70 specifically for the pathogen assays. For gene expression work and callose deposition assays infections were at CT26 and CT42. These times correspond to subjective morning and evening respectively.

2.3 Bacterial growth conditions

All bacteria were grown in King's Broth [159] supplemented with the required antibiotics (Table1). Bacteria were cultured at 30°C for 12 to 14 hours, corresponding to middle to late log-phase and prepared for inoculation as recommended in [160]. *P. syringae* DC3000 expresses avirulence B (avrB), an effector protein that is recognised by the R protein RPM1. *P. syringae* DC3000 expresses avirulence Rpt2, an effector protein recognised by the R protein RPS2. *P. syringae* *hrpA* lacks a TTSS and is therefore unable to deliver effectors.

Table 1. The various strains of *P. syringae* used and the required antibiotics

<i>Pseudomonas syringae</i> DC3000	Rifampicin 50µgmL ⁻¹
<i>Pseudomonas syringae</i> DC3000 avrB	Rifampicin 50µgmL ⁻¹ Tetracycline 15 µgmL ⁻¹
<i>Pseudomonas syringae</i> DC3000 avrRpt2	Rifampicin 50µgmL ⁻¹ Kanamycin 50µgmL ⁻¹
<i>Pseudomonas syringae</i> DC3000 <i>hrpA</i>	Rifampicin 50µgmL ⁻¹ Kanamycin 50µgmL ⁻¹

2.4 Pressure infiltrating *Arabidopsis* plants for pathogen assays

Plants were inoculated by pressure infiltration using a 1mL needleless syringe to inject the abaxial side of the leaf as described in [160]. Bacteria were pressure infiltrated into three leaves per plant with three biological replicates. Mock treated plants were infected with 10mM MgCl₂. After infection, the infected leaves were harvested to measure bacterial growth. This was done at 4 and 48h after infection. To measure bacterial growth, leaf discs were homogenised in 1mL of 10mM MgCl₂ and appropriate dilutions plated onto KB agar supplemented with the necessary antibiotics. Petri dishes were incubated in the dark for two days at 30°C. All infiltrations were under constant light with the plants remaining in constant light for the remainder of the experiment. Counts measured as colony forming units per cm² of leaf.

2.5 Callose deposition assays

Callose levels were compared in wild type *Arabidopsis* Col-0 (hereafter referred to as Col-0) and the overexpressing line, *CCA1-ox* at CT26 and CT42. Four-week-old plants were infected with *P. syringae* DC3000 *hrpA* mutant lacking type three-secretion system and callose levels were measured 14h after infection. Plants were infected with a bacterial suspension at an OD of 0.2 which corresponds to 1×10^8 cfu ml⁻¹ as recommended by [161]. Using a 1 mL needleless syringe, bacteria were pressure inoculated through the abaxial side of the plant leaf surface [161]. After 14 hours, the infected leaves were detached and subjected to staining by aniline blue [161]. Infections were under constant light and temperature conditions at two time points, CT26 and CT42. Mock-treated plants were pressure inoculated with 10mM MgCl₂ and also subjected to staining by aniline blue [161]. After staining, leaves were fixed on glass slides and sealed with nail polish. Images were collected using a fluorescent inverted microscope (Nikon TMD-EF DIAPHOT-TMD, Tokyo, Japan) of callose deposition and these were quantified using Quantity One software (Bio-Rad Laboratories, Hercules, California).

2.6 Measuring gene expression

2.6.1 Gene expression using luciferase

Promoter activity of *Pathogenesis Related1::LUCIFERASE* (*PR1::LUC*), *Oxidative Inducing Signal 1::LUC* (*OX1::LUC*) and *Cold and Circadian Regulated 2::LUC* (*CCR2::LUC*) was measured. Bioluminescence was measured in four-week-old plants inoculated with *P. syringae* DC3000 or mock treated with 10mM MgCl₂, using a Xenogen Lumina IVIS (Caliper Life Sciences, MA, USA) with thermoelectrically cooled, low-light-detecting charge-coupled device (CCD) camera, in a temperature-controlled dark box. Plants were sprayed with 5 mM luciferin (Biosynth AG, Switzerland) 24 h and 12 h before the start of experiment, and were subsequently sprayed every 8 h with 1mM luciferin during the experiment. Bioluminescence was captured for 5 min, every 2 h in the Lumina IVIS. In order to calculate levels of bioluminescence, Living Image v4.1 software (Caliper Life Sciences, MA, USA) was used. Total bioluminescence counts were divided by the area to yield counts per cm². Only infected leaves were measured and luciferase levels were measured in four biological replicates per time point. Plants were grown in 16h light and 8h dark cycles and transferred to constant conditions 24h prior to infection. Bioluminescence measurements were carried out 24h after infection at CT26 and CT42.

2.6.2 Quantitative PCR

2.6.2.1 RNA extraction and cDNA synthesis

Col-0 and *CCA1-ox* plants were grown for four weeks in 16h light/ 8h dark cycles, and were transferred to constant light at least 24 h before infection. Three plant leaves per plant of Col-0 and *CCA1-ox* were inoculated at CT26 and CT42 using a 1 mL needleless syringe with either *P. syringae* DC3000 or 10mM MgCl₂. Inoculated leaves were harvested in liquid nitrogen 4h after infection and stored at -80°C. RNA was extracted as described by Smart and Roden [162]. Quantity and integrity was tested by spectrophotometric readings (Nanodrop ND11) and separation on a 1.2% agarose gel by electrophoresis, respectively. For cDNA synthesis RNA was reverse transcribed using ImpromII reverse transcription system (Promega, Madison WI, USA), following the manufacturer's protocol except half reaction-volumes were used and random hexamer primers (Promega) were added. Two technical repeats for cDNA synthesis were carried out to account for variation in efficiency of cDNA synthesis. The pooled cDNA was used for quantitative-Polymerase Chain Reaction (qPCR) experiments. Three biological replicates were used for each treatment.

2.6.2.2 qPCR using standard curve method

Real Time PCR was carried out using a Rotor Gene 6000 machine (Corbett Life Science Pty. Ltd., Sydney, Australia). Transcript levels of *WRKY22*, *WRKY29* and *SRFR1* were measured in Col-0 and

CCA1-ox infected at CT26 and CT42. Basal expression was determined for *SRFR1* at CT26 and CT42 while basal and induced transcript levels were measured for *WRKY22* and *WRKY29* at CT26 and CT42. The Dynamo flash SYBR Green qPCR kit (Finnzymes, Keilaranta, Finland) manufacturer's protocol was followed except half reaction volumes were used. A total reaction volume of 10µl per reaction consisted of 1µl of cDNA template, 0.5 µl each of forward and reverse primers, 5µl of SYBR green mix and 3µl of sterile H₂O. The housekeeping gene was *UBIQUITIN10* (*UBIQ10*) and the genes investigated were *WRKY22*, *WRKY29* and *SRFR1*. *UBIQ10* was chosen as the housekeeping gene as it is expressed constitutively and is unaffected by the circadian clock [163]. Primers, annealing conditions and R² values are listed in Table 2. The R² values are above 0.995 for the housekeeping gene and *WRKY22* and *WRKY29*. *SRFR1* has a R² value of 0.988 (Table 1.). R² values of 0.995 indicate a linear standard curve. To determine copy number the standard curve method was used [164,165]. A standard curve was made, ranging from 1×10¹ copy numbers/µL to 1×10⁶ copy numbers/µL using purified PCR product of known concentration and base pair length for each gene investigated including the housekeeping gene. The R² values shown in Table 2 indicate excellent correlation coefficients for the standard curves. Using Rotor Gene 6000 software, copy number for the housekeeping gene was determined from the standard curve. Similarly, copy number for the genes of interest was also determined. To calculate relative gene expression the copy number of the gene of interest was divided by the copy number of the housekeeping gene.

Relative transcript abundance = gene of interest (copy number) ÷ housekeeping gene (copy number)

Table 2. Primers used, annealing conditions and R² values.

Primer	Primer sequence	Annealing temperature	R ² value
<i>UBIQ10f</i>	5' TTGTCGATGGTGTCCGAGCTT 3' [166]	Anneal at 60°C	0.997
<i>UBIQ10r</i>	5'TAAAAACTTTCTCTCAATTCTCTCT 3' [166]		
<i>WRKY22f</i>	5'CGACAAAGTAATGCCGTCTCC 3' [167]	Anneal at 55°C	0.996
<i>WRKY22r</i>	5'CGTTTCTGGTTCTGTGGCTTT 3' [167]		
<i>WRKY29f</i>	5'-ATCCAACGGATCAAGAGCTG-3' [168]	Anneal at 60°C	0.996
<i>WRKY29r</i>	5'-GCGTCCGACAACAGATTCTC-3' [168]		

<i>SRFRIf</i>	5' CTGG`ATATGCCTCACTAGAA3' [169]	Anneal at 60°C	0.988
<i>SRFRIr</i>	5' CACTGGGTCACAAGGCTCTG 3' [169]		

2.7 Online Bioinformatics tools

To obtain accession numbers for genes examined in this study the online *Arabidopsis* resource was used <http://www.arabidopsis.org/>. To analyse promoter regions of plant genes the online tool found at this link <http://www.bioinformatics2.wsu.edu/cgi-bin/Athena/cgi/home.pl> (*Arabidopsis thaliana* expression network analysis) (ATHENA) was used.

2.8 Statistical analysis

To determine whether time of infection and/or plant genotype affect bacterial titre and callose deposition general linear model (GLM) analysis was performed using Statistica v10.1 followed by the Bonferroni post-hoc test. To analyse bioluminescence data for *PR1* and *OXII* expression Two Way Repeated Measures ANOVA was used. Student's t-test was used to analyse qRT-PCR data. A p-value of less than 0.05 was considered significant.

Chapter 3: Results

3.1 Investigating the MAMP-triggered immune response for circadian control

3.1.1 Investigating the effect of time-of-day on *Arabidopsis* response to *P. syringae* DC3000:

To investigate the potential circadian regulation of plant defence responses, 4-week-old Col-0 plants were infected with the plant pathogen *P. syringae* DC3000 at 4h intervals under constant light and temperature conditions. As infections were carried out in free running conditions, the time at which plants were infected was recorded as circadian time (CT), i.e., the plant's subjective or internal time, shown on the x-axis (Fig 3.1). Subjective evening is indicated by the shaded regions in the figure (Fig 3.1).

Bacterial titres measured 48h after infection indicate that Col-0 plants were least susceptible to infection at CT26 and CT54, while greatest success of infection was recorded at CT42 and CT66. Generally, the trend suggested infections during subjective day led to a better defence response than infections during the subjective night, with lower bacterial counts recorded for the former (Fig 3.1). This time-of-day difference in susceptibility to infection is observed through two circadian cycles. The peaks and nadirs in susceptibility to infection under free-running conditions indicated a possible circadian clock driven influence on plant responses to *P. syringae* DC3000. To investigate this possibility, defence responses of Col-0 plants were compared with defence responses in clock deficient plants under free running conditions.




Figure 3.1 *Arabidopsis thaliana* (Col-0) shows variable susceptibility to *P. syringae* DC3000 under free-running conditions of constant light. Bacterial growth in leaves was measured 48h after infection under constant light conditions at each of the 12 time points. The success of growth was used as a measure of virulence (measured as colony forming units per cm^2 cfu cm^2). The error bars indicate standard error of mean for three measurements from three biological replicates. A similar pattern in the variation in susceptibility was observed in three independent experiments. (Absence of error bars implies little variation).

3.1.2 Investigating *elf3-1* and *CCA1-ox* for time-of-day variation in susceptibility to *P. syringae* DC3000:

The circadian clock null mutant, *elf3-1* and *CCA1*-overexpressing line *CCA1-ox*, were investigated for time-of-day variation in susceptibility to infection by *P. syringae* DC3000 (Fig 3.2). Under constant light conditions *CCA1-ox* has been shown to have decreased fitness, arrhythmic circadian regulation and lower photosynthesis [101], while in constant light, the null mutant *elf3-1* was reported to be arrhythmic [170]. Recently researchers showed that the *cca1* null mutant displayed no time-of-day variation in susceptibility to *Hyaloperonospora arabidopsis* (*Hpa*) [146] when infected under light/dark cycles. In addition, *CCA1-ox* showed enhanced disease resistance to *Hpa* [146] compared to Col-0 after infections in the morning.

In this study, plants were infected under free-running conditions of constant light. Col-0, *elf3-1* and *CCA1-ox* were infected at CT26 and CT42 as these time points corresponded to least susceptible and

most susceptible in Col-0, respectively (Fig 3.1). Bacterial titres at 4hpi indicated that equal numbers of bacteria were inoculated across time points and genotypes (Fig 3.2). Plant susceptibility to *P. syringae* DC3000 was measured after 48h as in Fig 3.1. Once again, Col-0 showed time-of-day variation to pathogen infection with lower titres recorded at CT26 than at CT42 (Fig 3.2). Interestingly, the arrhythmic plants displayed equal susceptibility to infection, regardless of when the infection was carried out. Furthermore, these arrhythmic plants showed greater susceptibility to infection than Col-0 plants at CT26 with bacterial titres comparable to Col-0 at CT42. To test whether these differences were statistically significant, a general linear model (GLM) analysis comparing the effects of genotype and time was used followed by Bonferroni's post-hoc test grouping bacterial counts based on homogeneity. Similar counts were assigned the same letter indicating no significant difference in mean value (Fig 3.2).

The statistical analysis revealed that Col-0 plants showed a time-of-day variation in susceptibility to infection with bacterial levels significantly lower at CT26 compared to CT42 ($p < 0.05$). However, this time-of-day variation was absent in the arrhythmic plants, *elf3-1* and *CCA1-ox* (Fig 3.2). Collectively these results indicated that Col-0 showed the least susceptibility to infection at CT26 and displayed a time-of-day variation in defence response while the arrhythmic plants, *elf3-1* and *CCA1-ox* showed no time-of-day variation to infection and higher bacterial counts comparable to Col-0 at CT42. This further demonstrated a clock controlled time-of-day variation in defence responses of Col-0 but not of the arrhythmic plants.




Figure 3.2 Arrhythmic plants show no variation in susceptibility with respect to time of infection under free-running conditions of constant light. The wild type, *elf3-1* null mutant and *CCA1-ox* were infected with *P. syringae* DC3000 at CT26 and CT42. Bacterial growth at 48h after infection shows variation in susceptibility with time of infection in Col-0 but not in *CCA1-ox* and *elf3-1*. To test whether time of infection and genotype were significant, a general linear model was used followed by Bonferroni's post hoc test ($p < 0.05$). Dissimilar letters above the bars represent statistically significant differences in mean bacterial counts. Each measurement is the average of three biological repeats and error bars indicate standard error of mean (Note small error indicated little variation).

3.1.3 Investigating time-of-day variation in *CCA1-ox* in the second cycle:

To confirm observations made in Fig 3.2 that Col-0 showed a time-of-day variation in susceptibility but not *CCA1-ox*, Col-0 and *CCA1-ox* were infected in the second cycle at CT50 and CT66 that correspond to subjective morning and subjective evening infections respectively (Fig 3.3). Bacterial titres were measured at 4h post infection (hpi) and 48hpi. The 4hpi counts show equal numbers of bacteria inoculated at both time points and across plant genotypes. At 48h after infection, bacterial counts are lower for Col-0 at CT50 than at CT66 while *CCA1-ox* shows levels comparable to Col-0 at CT66 for both infection times. To test whether Col-0 levels were significantly lower at CT26, a two-tailed Student's t-test was carried out, giving a p-value of 0.037. This indicates that Col-0 is significantly lower at CT20. Additionally a p-value of 0.2852 after a two-tailed Student's t-test showed that mean counts in *CCA1-ox* at CT50 and CT66 are not significantly different.

These infections revealed that time-of-day variation to *P. syringae* DC3000 occurs in Col-0 plants while arrhythmic plants show equal susceptibility to infection. Furthermore, plants lacking a functional clock also show a greater susceptibility to infection than Col-0 at CT50, with levels comparable to Col-0 at CT66 (Fig 3.3).




Figure 3.3 Second cycle infections show time-of-day variation in Col-0 but not in CCA1ox. Levels of bacteria 48h after infections in Col-0 are lowest at CT50 while at CT66 are comparable to CCA1-ox mutant. Error bars represent standard error of mean from three biological replicates. A two tailed Student's t-test showed Col-0 at CT50 was significantly lower than at CT66 (shown by asterisk, $p < 0.05$). No significant differences were reported in CCA1-ox.

3.2 Investigating the ETI response of *Arabidopsis* for time-of-day variation under free running conditions:

3.2.1 Infecting wild type and clock mutants with *P. syringae* DC3000 avrB and avrRpt2 at CT26 and CT42:

Next, the effect of time-of-day variation on ETI was investigated using two avirulent strains: *P. syringae* DC3000 avrB and *P. syringae* DC3000 avrRpt2. These strains possess effectors that are recognised by the plants cognate R proteins thus initiating ETI.

P. syringae DC3000 avrB initiates ETI if the infected plant possesses the cognate resistance (R) protein RPM1 [171]. Absence of RPM1 results in successful infection with avrB enhancing virulence [171]. Col-0, *CCA1-ox* and *elf3-1* were infected with *P. syringae* DC3000 avrB at CT26 and CT42 and bacterial counts carried out 4h and 48h after infection (Fig 4i). Titres at 4h showed equal numbers of bacteria infected across time points and plant lines (Fig 4i). Bacterial counts at 48hpi indicated levels of bacteria were lowest in Col-0 at CT26, with slightly higher counts at CT42 (Fig 4i). Both *CCA1-ox* and *elf3-1* showed higher bacterial titres at CT26 than CT42 with *elf3-1* showing greater susceptibility to infection than Col-0 at both time points (Fig 4i). To examine if these differences were statistically significant GLM analysis and Bonferroni post-hoc analysis were performed. These tests suggested the observed differences in Fig 4i are not statistically significant with $p > 0.05$.

P. syringae DC3000 avrRpt2 activates ETI if the plant possesses the cognate R protein RPS2 [171]. If the plant does not have the R protein, avrRpt2 enhances infection. To investigate the effect of time-of-day, Col-0, *CCA1-ox* and *elf3-1* were infected with *P. syringae* DC3000 avrRpt2 (Fig 4ii). Pathogen assays at 4hpi show equal bacterial titres across genotypes and plant lines (Fig 4ii). After 48h, the lowest counts were noted at CT26 in Col-0, with slightly higher counts at CT42 in Col-0. Bacterial levels in *elf3-1* at both CT26 and CT42 were comparable to Col-0 at CT42 while *CCA1-ox* was the most susceptible with levels higher than both Col-0 and *elf3-1* (Fig 4ii). To investigate whether these differences were statistically significant GLM analysis followed by Bonferroni post hoc analysis were carried out. Interestingly GLM analysis suggested the effect of genotype alone on bacterial counts was significant ($p = 0.025$) and Bonferroni post-hoc test identified Col-0 as behaving differently from *CCA1-ox* but similar to *elf3-1* ($p = 0.0395$). However, when investigating the effect of genotype and time Bonferroni post-hoc test showed no significant differences, which suggests no effect of the clock on ETI response.




Figure 4 Bacterial counts in *Col-0*, *CCA1-ox* and *elf3-1* after 4 and 48h when infected with (i) *P. syringae* DC3000 *avrB* and (ii) *P. syringae* DC3000 *avrRpt2* at CT26 and CT42. Error bars represent standard error of mean for three biological replicates. Experiment performed only once.

3.3 Measuring callose levels of *Arabidopsis* in response to *P. syringae* *hrpA* at CT26 and CT42:

Fig 3.1 shows time-of-day variation in *Col-0* plants to infection with *P. syringae* DC3000 suggesting the circadian clock perhaps modulates MTI. One of the early responses to flg22 and non-pathogenic *P. syringae* is the production of callose regulated by callose synthase (*GSL5/PMR4*) [54]. *In silico* analysis of *PMR4* (At4g03550) using the online bioinformatics tool (ATHENA) found at this link- <http://www.bioinformatics2.wsu.edu/cgi-bin/Athena/cgi/home.pl>, showed *GSL5/PMR4* to have a CCA1 element in the promoter region (-405 to -398bp upstream of +1). The presence of circadian clock binding motifs in *PMR4* and the earlier result reported in Fig 3.1 indicated that the circadian clock may modulate callose production in a time-of-day manner. To investigate this, *Col-0* and *CCA1-ox* were infected with non-pathogenic *P. syringae* DC3000 *hrpA* at CT26 and CT42 under free running conditions (Fig 5.1 and 5.2). As *P. syringae* *hrpA* lacks a functional type III secretion system, it is unable to deliver effectors, and so is non-pathogenic [172]. Furthermore, detection of MAMPs from *P. syringae* *hrpA* initiates a strong MTI response. Callose levels were measured 14h after infection at either CT26 or CT42 (Fig 5.1 and 5.2). The highest callose levels were recorded at CT26 in *Col-0* while much lower callose levels were detected in *Col-0* at CT42 (Fig 5.1) Callose levels of *CCA1-ox* were greatly reduced at both time points with levels comparable to those of *Col-0* at CT42 (Fig 5.1). These results mirror observations made in Fig 3.1 and Fig 3.2.

To test if the observations in Fig 5.1 were significantly different, GLM analysis testing the effect of genotype and time of infection and Bonferroni post-hoc analysis were performed. GLM analysis showed both time and genotype ($p=0.00002$ and $p=0.02$ respectively) to be significant when factored

independently. Furthermore interaction between genotype and time was highly significant ($p=0.0006$). Callose levels with similar counts were grouped together using the Bonferroni post-hoc analysis. Callose counts showing significantly different levels ($p<0.05$) are represented with different letters (Fig 5.1). Notably, callose deposition showed time-of-day variation in Col-0 but not in *CCA1-ox*.




Figure 5.1 Callose levels measured at CT26 and CT42 in Col-0 and *CCA1-ox*. Levels of callose measured 14h after infection with *P. syringae hrpA* show higher production of callose at CT26 in Col-0 than at CT42. Production of callose is abrogated in *CCA1-ox* with levels comparable to Col-0 at CT42. Error bars are standard error of mean from six biological replicates. Letters above each bar represent statistically significant levels of callose (GLM analysis, Bonferroni's post-hoc test $p<0.05$). Experiments were performed twice with similar results.




Figure 5.2 Fluorescent microscope images showing callose deposits. Callose deposits appear as white specks when they fluoresce. For each plant, two leaves were infected and stained with aniline blue 14h later. Several regions of the infected leaves were photographed and quantified. (Scale bar = 0.2mm).

3.4 Promoter activity and expression of genes involved in the MTI response to *P. syringae* DC3000 infection:

Given that the circadian clock plays a role in plant defence [146], and these experiments suggested MTI could be modulated by the clock, investigating genes involved in MTI was essential. Upon recognition of MAMPs, one of the immediate results is the production of active oxygen species (AOS) such as H₂O₂ [44]. This AOS burst activates Oxidative Signal Inducing 1 (OXI1) which is important for full MTI response to *P. syringae* [43]. OXI1 is also necessary for the full activation of the mitogen activated protein kinases, MPK3 and MPK6 [44]. In turn, the MAP kinase cascade involving MPK3 and MPK6 is known to be up regulated in response to MAMPs, activating *WRKY22* and *WRKY29* which are important downstream components of the MAPK cascade [45]. The *WRKY* genes are a plant-specific group of transcription factors and are important in regulating defence gene expression upon infection [45,167,173]. One of the outputs of this pathway is the expression of *PATHOGENESIS RELATED 1 (PR1)*. Importantly though *PR1* is not an early response gene, occurring much later in the defence response [45] and is expressed in response to accumulation of SA [28]. *PR1* is an important defence signal in response to biotrophic pathogens such as *P. syringae* and is important in SAR [174,175]. Here promoter activity of *OXI1* and *PR1* was investigated in response to infection with *P. syringae* DC3000. Additionally, mRNA induction of *WRKY22* and *WRKY29* genes were measured 4h after infection.

3.4.1 Promoter activity of *OXI1* following infection

Wild type plants with the *OXI* promoter fused to a luciferase construct were infected with *P. syringae* DC3000. *OXI1* promoter activity was measured as bioluminescence in response to infection. An upstream analysis of the promoter region of *OXI1* indicated the presence of two putative CCA1 binding elements (-580 to -573 and -949 to -942 from start). Plants were infected at CT26 and CT42 and levels of bioluminescence measured from 24h after infection (Fig 6.1). Bioluminescence levels and peak expression levels for both infection time points were comparable with no significant differences reported after a Two-Way Repeated Measures (RM) ANOVA.




Figure 6.1 *OX1* expression measured at CT26 and CT42 from 24h after infection with *P. syringae* DC3000. Error bars represent standard error of mean when four biological replicates were infected per time point. A 2 way RM ANOVA showed no significant differences in *OX1* expression profiles at CT26 and CT42 ($p < 0.05$). A similar result was observed in an independent experiment.

3.4.2 Promoter activity of *PR1* following infection

PR1 promoter activity was also investigated using a luciferase reporter construct. Wild-type *PR1::LUC* plants were infected with *P. syringae* DC3000 under free-running conditions at CT26 and CT42 and bioluminescence was measured from 24h after infection (Fig 6.2).

There is quicker induction of *PR1* following infection at CT42 than at CT26 based on promoter activity of *PR1* measured as bioluminescence (Fig 6.2). A Two Way RM ANOVA confirms this difference in induction of *PR1* as statistically significant ($p < 0.001$). Levels of *PR1* are higher at CT42 than at CT26 from 40h after infection, remaining higher until 56h after infection where levels begin to overlap.

PR1 expression occurs much later and is associated with SAR in uninfected leaves, HR, and PCD in infected leaves. While callose experiments (Fig 5.1 and 5.2) indicated a stronger response during subjective morning, complementing earlier observations (Fig 3.1 and 3.2) paradoxically, *PR1* levels were higher after infection at subjective night. We next investigated *WRKY22* and *WRKY29* which encode transcription factors important in activating early defence response genes and are themselves rapidly induced [45].




Figure 6.2 shows *PR1* expression measured as bioluminescence at CT26 and CT42. *PR1::LUC* Col-0 plants infected with *P. syringae* DC3000 at CT26 and CT42 are shown with open and closed circles respectively. Error bars are for four biological replicates and signify standard error of mean. A similar result was observed in an independent experiment. The asterisks indicate a significant difference in induction between at CT42 and CT26 ($p < 0.001$, 2 Way RM ANOVA).

3.4.3 Measuring transcript levels of *WRKY* transcription factors

WRKY22 and *WRKY29* are downstream of the MAPK pathway and are important in early defence responses to infection [45]. Activation of *WRKY22/29* results in activation of defence genes. Col-0 and *CCA1-ox* plants were infected at CT26 and CT42 with either 10mM $MgCl_2$ or *P. syringae* DC3000, (mock and treated, respectively in Fig 6.3). Expression levels of *WRKY22* (Fig 6.3i) and *WRKY29* (Fig 6.3ii) were measured 4h after inoculation in mock and treated plants.

Expression of *WRKY22* was induced in both genotypes at both time points (Fig 6.3i). The mock treated plants show un-induced levels of *WRKY22* which corresponds to basal levels. These basal levels are similar across both genotypes and time points. While induction of *WRKY22* appears variable, the large error bars suggest this difference is not significant (Fig 6.3i). A Student's t-test confirmed that no significant differences could be observed in induced and basal levels of *WRKY22* across both genotypes at CT26 and CT42 (Fig 6.3i).

WRKY29 possesses both CCA1 and EE binding motifs in its promoter region. A CCA1 motif Binding Site in Chlorophyll A/B Binding protein 1 (CCA1 motif BS CAB1) is also reported within the CCA1 motif. The position of the CCA1 motif is from -151 to -144 base pairs upstream of the start site while the CCA1 motif BS in CAB1 is from -152 to -143 base pairs from the start site. The evening element is located -314 bases from the start site.

Basal levels of *WRKY29* were similar across genotypes and time points. Though levels of *WRKY29* appear higher in mock-infected *CCA1-ox* at CT26 compared to CT42, the large error associated with this measurement meant no significant differences in variation were observed. Additionally, induced *WRKY29* levels in *CCA1-ox* were similar at both CT26 and CT42. A Student's t-test comparing induced levels of *WRKY29* in *CCA1-ox* showed no significant differences between the two time points ($p = 0.2663$) (Fig 6.3ii). However, *WRKY29* levels in Col-0 showed higher levels in plants infected at CT26 than at CT42 (Fig 6.3ii). A two-tailed Student's t-test revealed a statistically significant difference ($p = 0.0105$) suggesting significantly higher levels of *WRKY29* are induced at CT26 than at CT42. As basal levels of *WRKY29* in Col-0 are similar, this is potentially a very interesting observation; but these findings are based on one experiment only and therefore must be interpreted with caution.




Figure 6.3 Gene expression levels of (i) *WRKY22* and (ii) *WRKY29*. Mock and treated plants were inoculated with 10mM $MgCl_2$ and *P. syringae* DC3000 respectively. Shaded bars represent *CCA1-ox* while un-shaded bars represent Col-0. The plants were infected at CT26 and CT42, tissue harvested 4h later and transcript levels measured by qPCR. Error bars represent standard error of mean for three biological replicates. A 2 tailed Student's t-test comparing induced levels of *WRKY29* show levels are higher at CT26 compared to CT42 ($p < 0.05$) as shown by the asterisk. One experiment only.

3.4.4 *SRFR1* levels at CT26 and CT42

Wang *et al.* (2011) recently showed that plants regulate a number of defence genes to anticipate dawn infection [146]. The authors showed that defence genes are pulse expressed at dawn when likelihood of infection is greatest and that in the event of infection, prolonged expression of these defence genes results in disease resistance and sometimes PCD [146]. Using an avirulent oomycete, *Hpa Emwa1*, they show that plant defence genes time their expression to peak when *Hpa* releases its spores [146]. Furthermore, they showed the circadian clock modulates expression of the defence gene *RPP4*, with its expression peaking at dawn [146].

In a separate study, a *srfr1* null mutant showed overexpression of *RPP4* and enhanced disease resistance [87]. This study revealed *SUPPRESSOR OF RPS4-RLD1 (SRFR1)* as a negative regulator of NB-LRR proteins, preventing over-accumulation of R proteins thus preventing autoimmunity [87,176,177]. These independent observations led me to predict that *SRFR1* expression could be antagonistic to *RPP4*. *RPP4* has a circadian rhythm with peaks and nadirs at dawn and dusk respectively. I investigated whether *SRFR1* would follow a similar but antagonistic rhythm with peaks at night and nadirs during the day.

Basal levels of *SRFR1* were measured during subjective day and subjective night (CT26 and CT42 respectively) (Fig 6.4). Normalised expression of *SRFR1* shows higher levels of *SRFR1* at CT42 compared to CT26. A two tailed Student's t-test confirmed that these observations are statistically significant ($p=0.007$). However, replicate experiments are required, using a larger sample size to verify observations made in this preliminary experiment.




Figure 6.4 Basal levels of *SRFR* in Col-0 at CT26 and CT42. *SRFR1* transcript levels were measured from three biological replicates. Expression of *SRFR1* was normalised using Col-0 CT26 levels. A Two-tailed Student's t-test comparing basal levels of *SRFR1* at CT26 compared to CT42 showed levels of *SRFR1* are higher at CT42 ($p=0.007$, represented by double asterisks). One experiment only.

3.4.5 Investigating circadian clock function during infection

AtGRP7 also known as *COLD AND CIRCADIAN REGULATED (CCR2)* is an evening-phased gene [178] under the control of the circadian clock [166,179,180,181]. Wang *et al.* (2011) used *CCA1* and *LHY* promoter-luciferase reporter constructs, to measure promoter activity in free-running constant light conditions after infection with *Hpa* Emwa1 [146]. Infection resulted in significant induction and arrhythmic expression of *CCA1* while similar measurements of *LHY* indicated that it was unaffected [146]. As *CCA1* is an important component of the circadian clock, this could lead to disruption of the circadian clock or outputs of the clock because of infection. To investigate this, *CCR2* promoter activity was measured before and during infection. *CCR2::LUC* showed a circadian rhythm, with levels peaking in the evening and this rhythmic expression was unaffected after infection (Fig 6.5).

The results indicate core clock function is not disrupted by infection with *P. syringae* (Fig 6.5) complementing work by Wang *et al.* (2011) showing overall clock function is not disrupted after infection of *Arabidopsis* with *Hpa* under free running conditions [146].




Figure 6.5 Expression profiles of *CCR2::LUC* before and after inoculation with mock (10mM MgCl₂) or infected (*P. syringae* DC3000) under free running conditions (LL). Inoculation was at 24h, indicated by arrow. Error bars are standard error of mean for four biological replicates.

Chapter 4 Discussion

The results reported here indicate that Col-0 plants display time-of-day variation in susceptibility to *P. syringae* DC3000 (Fig 3.1). While previous reports have documented time-of-day variation to plant-pathogens they attributed it to the role of light [182] and the circadian clock [146]. Both these observations were made under light/ dark cycles but to implicate one or the other it was necessary to carry out the infections under free-running conditions [10,157]. The findings in this report indicate that success of infection varies with time-of-day with lowest infection at subjective morning (CT26 and CT50) and greatest infection at subjective night (CT42 and CT66) (Fig 3.1). Success of infection was used as a measure of strength of plant defence response, suggesting that plant defence is better primed for defence at CT26/50 than at CT42/66 (Fig 3.1). Plants activate MTI upon infection by *P. syringae* DC3000 and the reported results suggest that the circadian clock modulates MTI. A stronger MTI response is observed after infection during subjective day than subjective night. While the time-of-day differences in susceptibility in Fig 3.1 are similar to those made under light/dark cycles [146] the data in Fig 3.1 is obtained from infections performed under free-running conditions. The persistence of time-of-day variation in strength of defence responses under free-running conditions firmly implicates control of MTI by the circadian clock.

Infecting the clock mutant *elf3-1* and *CCA1-ox* revealed them to have diminished defence responses and to be equally susceptible to infection in the morning and evening, unlike Col-0 plants (Fig 3.2 and Fig 3.3). This result complements similar findings using a *cca1* null mutant after infection under light-dark cycles with *Hpa* [146]. Contrastingly, *CCA1-ox* was shown to have enhanced defence responses to *Hpa* when infected under light dark cycles [146]. Findings reported here indicate that *CCA1-ox* showed diminished levels of defence with bacterial titres higher than Col-0 at CT26 but comparable to Col-0 at CT42 (Fig 3.2). Therefore, this result contradicted previous work by Wang and colleagues (2011) but there were notable experimental differences between the two studies. They used an oomycete pathogen and performed infections under alternating light-dark cycles while experiments in this report used a bacterial pathogen and were performed under free-running conditions of constant light. One could speculate that under light/dark cycles where there is some degree of driven rhythmicity [183], the over-expression of *CCA1* confers an advantage of heightened disease resistance while under free-running conditions this advantage is lost. Previous work reported that under alternating light dark cycles a number of physiological processes retain rhythmicity in the *CCA1-ox* mutant but are lost upon transfer to free running conditions [183].

One of the classic MAMP triggered responses (MTI) is the production of callose. *In silico* analysis, revealed *GSL5/PMR4* encoding a callose synthase possesses a CCA1 binding motif (found -405 to -398 from +1) in its promoter region. These promoter motifs are small and can occur by chance thus their presence alone is not sufficient to attribute regulation by a certain component. Nonetheless, the presence of these motifs warranted further investigations into whether callose production may be regulated by the circadian clock. Callose levels were measured 14h after infection at CT26 and CT42 under free running conditions in Col-0 and *CCA1-ox* (Fig 5.1 and Fig 5.2). Our results indicate that callose production is seven times greater at CT26 than at CT42 in Col-0. Additionally *CCA1-ox* displayed abrogated levels of callose, comparable to Col-0 at CT42 and showed no significant time-of-day variation (Fig 5.1). These results complemented observations in Fig 3.2 where Col-0 showed time-of-day variation in susceptibility to infection while *CCA1-ox* did not. Collectively these findings indicated that the circadian clock modulates the MTI response and that callose response, a classical marker of MTI varied in strength depending on when the plant was infected. This time-of-day variation of the MTI response and callose production in response to infection by an equal quantity of bacteria at different times suggests that these defence responses may be gated by the circadian clock.

PR1 expression is mediated by the salicylic acid signalling pathway and is part of the systemic response as well as the MTI and ETI responses [84]. Using a luciferase reporter construct, *PR1* promoter activity was measured (Fig 6.2). Paradoxically, *PR1* showed quicker induction after infection at subjective night (CT42) compared to subjective morning (CT26). This result contradicted other results that had indicated a strong defence response during the day. A recent study by Sato and colleagues (2010) provided an intriguing explanation of events. They showed that callose synthase and the salicylic signalling sector mutually inhibited each other [54]. By dividing plant defence genes into sectors based on positive regulatory relationships they identify an early MAMP triggered (EMT) sector of which callose synthase is a member and salicylic acid sector [54]. They showed that these two sectors are mutually antagonistic and were capable of inhibiting each other thus when the EMT sector was fully activated it negatively regulated the salicylic acid sector and *vice versa* [54]. My results indicated a strong EMT response at CT26 and a weak EMT response at CT42 as measured by callose production (Fig 5.1). Based on the observations made by Sato and colleagues, one could speculate that in this study the strong EMT observed at CT26 in Col-0 would mutually inhibit the SA sector thus delaying induction of *PR1*. Furthermore, the weak EMT response observed at CT42 in Col-0 (Fig 5.2) would mean the associated antagonistic effect on the SA sector is weaker at CT42 thus *PR1* accumulated quicker at CT42. These *PR1* data support this hypothesis. Additionally, another early response gene, *WRKY29*, showed greater induction at CT26 than at CT42 (Fig 6.3) lending further support to the proposed idea. Collectively, one could surmise that possibly the time-of-day variation in *PR1* was a downstream effect of varying strength of early defence response genes.

Transcript levels of *WRKY22* and *WRKY29* were also investigated as they are both induced in response to pathogens [45,167,173]. Preliminary data reported here suggested that the circadian clock may gate the strength of *WRKY29* response to bacteria (Fig 6.3ii). A Student's two-tailed t-test showed that *WRKY29* levels were greater at CT26 than at CT42 in infected plants. Mock treated plants showed similar *WRKY29* basal levels at both CT26 and CT42 in Col-0. This indicated that the amplitude of response in Col-0 was greater at CT26 than at CT42. This data, though promising is acknowledged as preliminary data. Furthermore, the biological contribution of such small differences though statistically significant must be verified. Importantly, levels of *WRKY29* in infected *CCA1-ox* were comparable to those of infected Col-0 levels at CT26 (Fig 6.3ii) and showed no time-of-day variation. The absence of time-of-day variation in levels of *WRKY29* in *CCA1-ox* complements other findings in this study (Fig 3.2, Fig 3.3 and Fig 5.1) however; transcript levels are similar to wild type at CT26, which does not explain the higher susceptibility to infection of *CCA1-ox* (Fig 3.2). Thus, while expression of *WRKY29* showed time-of-day variation in Col-0 and not in *CCA1-ox*, expression levels in the arrhythmic mutant were comparable to wild type levels. This weakens attempts to draw significant biological conclusions but warrants further investigations.

In addition, *WRKY22*, which is functionally redundant to *WRKY29*, behaved similarly across time points and genotypes, showing similar basal levels and similar amplitude of induction (Fig 6.3 ii). Collectively then, the *WRKY* dataset suggests that *WRKY* genes investigated here are equally inducible at either night or day in both Col-0 and *CCA1-ox* and therefore they are unlikely to explain the time-of-day variation observed in previous results. While levels of *WRKY29* in Col-0 plants showed statistically significant differences at different times, this result is based on three biological replicates and should be repeated in independent experiments to verify robustness and reproducibility of this observation. The *WRKY* family of transcription factors positively and negatively regulate defence responses therefore other members of this family could be ideal candidates for further investigation.

Recent results point to plant defences peaking at dawn, showing a 'pulse' in expression of defence genes [146]. One such gene displaying pulse expression at dawn is *RPP4* [146], a defence gene involved in plant immune responses to infection. *RPP4* is also negatively regulated by *SRFR1* [87]. The observation that *RPP4* displayed pulse expression at dawn and that it is also negatively regulated led to the possibility that its negative regulator may show an antagonistic expression profile. Data reported here indicated that *SRFR1* is lower during the 'subjective' day than 'subjective night' (Fig 6.4). This result suggests that negative regulators of defence may themselves experience down regulation at dawn, thus facilitating induction of defence genes. Suppressors may follow an

antagonistic circadian rhythm, showing decreased levels at dawn and increased levels at night. While results reported here (Fig 6.4) supported this claim, the difference between levels of *SRFR1* at CT26 and CT42 was less than 25%. Whether such a difference in transcript levels is biologically significant requires further investigation. Additionally this observation was made based on one experiment and therefore must be replicated in larger numbers to ensure that this observation is both robust and replicable. Perhaps the more likely explanation has to account for both down-regulation of suppressor genes and up-regulation of defence genes that results in the pulse expression observed at dawn.

Most of the findings reported here indicate that the circadian clock may modulate MTI. Wang and colleagues showed that the circadian clock also regulated genes encoding R proteins [146]. To investigate whether defence responses showed time-of-day variation with specific R protein and effector protein interaction, Col-0, *CCA1-ox* and *elf3-1* plants were infected at CT26 and CT42. Infections were carried out with *P. syringae* DC3000 avrB and avrRpt2 (Fig 3.3i and ii).

Infections with *P. syringae* DC3000 avrB showed no significant time-of-day variation (Fig 3.3i). Statistical tests indicated that no significant differences could be observed across various infection times and genotypes. While these results indicated that defence responses to *P. syringae* DC3000 avrB showed no time-of-day variation, *P. syringae* DC3000 avrRpt2 showed some differences. Statistical analysis (GLM and Bonferroni post-hoc test) indicated that Col-0 behaved significantly differently from *CCA1-ox* but similar to *elf3-1*. From this data, one could surmise that certain R proteins involved in recognition of avrRpt2 may be affected in *CCA1-ox* plants affecting their ability to recognise and initiate defence. Alternatively, the variation between Col-0 and *CCA1-ox* may hint at a role for CCA1 in plant defence responses, drawing from findings by Wang *et al.* 2011. In contrast, time-of-day and the circadian clock might not affect the recognition mechanisms involved in activation of defence responses to avrB. These observations are mere speculations as the data available are based only on one experiment and further work is necessary to draw significant conclusions. As, Wang *et al.* 2011 reported a number of R defence genes to be under control of CCA1 further experiments should target those R protein interactions using cognate avirulent strains.

An early event upon MAMP perception is production of active oxygen species such as H₂O₂. The production of H₂O₂ results in the induction of *OXII* which encodes a serine/threonine kinase [43,44]. OXII kinase is activated by H₂O₂ and is essential for full activation of the mitogen-activated protein kinases MPK3 and MPK6 [44]. *In silico* analysis revealed that *OXII* has a CCA1 binding motif in its

promoter region, which suggested possibility for circadian regulation in the absence of pathogens. Promoter activity of *OXII* showed no variation in expression profiles between infections at CT26 and CT42 (Fig 6.1) however *OXII* promoter activity was measured 24h after infection (Fig 6.1). As *OXII* induction occurs after an AOS burst, which occurs within 30min of perception of MAMPs, measuring early *OXII* expression may give a better indication as to whether induction varies with time-of-day.

Lastly, this report showed that *CCR2* expression is unaffected after infection with *P. syringae* DC3000 (Fig 6.5). *CCR2* is an evening phased gene regulated by CCA1 and LHY [181] and has been implicated in regulation of stomata [155] and plant defence [156]. Global transcript profiling of an over-expresser of *CCR2* showed enrichment of abiotic and biotic stress response genes including *PR1* [136]. Notably a point mutation in the RNA binding domain of *CCR2* diminished any such enrichments showing that the regulation involves the RNA-binding activity of *CCR2* [136]. The role of *CCR2* in defence is underscored by the observation that it is a target of the bacterial effector HopU1 [156]. HopU1 enhances virulence by ribosylating the RNA binding domain of *CCR2* interfering with its ability to bind RNA [156]. The observation that *CCR2-ox* showed enrichment of *PR1* and that *CCR2* expression peaked in the evening (Fig 6.5) may offer an alternate explanation as to why *PR1* showed quicker induction after subjective night infections (Fig 6.2).

Wang *et al.* (2011) demonstrated that *CCA1* expression was rapidly induced and became arrhythmic after infection with *Hpa* [146]. They measured both *CCA1* and *LHY* expression and unlike *CCA1*, *LHY* retained rhythmic expression and was unaffected by pathogen attack [146]. Moreover, *cca1* null mutant plants showed attenuated defences while *lhy* mutants did not [146]. Collectively these results suggest that CCA1 plays a role in a defence output pathway regulated by the clock rather than via the central clock mechanism [184]. As LHY and CCA1 are functionally redundant, Wang and colleagues predicted that overall clock function was unaffected even though *CCA1* becomes arrhythmic [146]. The findings reported here show that *CCR2* retains rhythmic expression after infection with *P. syringae* DC3000 under free-running conditions. As *CCR2* expression is under the control of CCA1 and LHY this would suggest two possible explanations. The first is that functional redundancy of *CCA1* and *LHY* prevents the *CCR2* pathway from losing rhythmic expression after infection. Secondly, induction of *CCA1* may be unique to *Hpa* Emwa1 infection only. Therefore, *CCA1* levels should be measured in response to infection by *P. syringae* DC3000.

The results presented here as well as those reported by other researchers [146,150,151] demonstrated that the circadian clock modulates plant defence responses to *P. syringae* DC3000 with peak defence occurring in the morning and greatest susceptibility to infection at night. The plant innate immune system and the plant circadian clock are complex aspects of plant physiology and independently are the subject of constant investigation. Recent findings including data reported here show an interface of cross talk exists between the plant circadian clock and its defence responses. Why this is so is a crucial question. Wang *et al.* (2011) offer a possible answer. The circadian clock in plants enables them to anticipate regular diurnal events, priming physiological and biochemical processes to occur at the most advantageous time. Wang and colleagues (2011) showed that defence responses were ‘pulse’ expressed at dawn. Using *Hpa* because its infection stages were clearly defined [185] they showed that ‘pulse’ expression at dawn coincided to when *Hpa* released its spores. They concluded that the lifecycles for biotrophic pathogens are intimately linked with their host metabolism, which is controlled by the circadian clock.

P. syringae DC3000 is also a biotrophic pathogen and showed time-of-day variation in its ability to infect *Arabidopsis* plants (Fig 3.1). While numerous researchers have documented various aspects of *P. syringae* DC3000, a clear understanding of the stages of infection has not been completely developed. A possible model of infection could be that plants open and close their stomata with a circadian rhythm, opening stomata at dawn and closing them at night. Stomata are important for gaseous exchange, transpiration and photosynthesis and are a natural portal of entry that bacteria use. This would suggest the ideal time for bacteria to infect plants would be when stomata are fully open, at dawn, and conversely plants anticipate this and so up regulate defence genes to minimise infection. Results reported in Fig 3.1 were after pressure infiltration, thereby bypassing plant stomata altogether and perturbing defence responses at both subjective day and subjective night. A stronger defence response was observed by lower counts at subjective day and a weaker defence response at subjective night (Fig 3.1). This result hints at the possibility that at night when stomata are shut, the anticipated risk of pathogen entry is less and so defence responses are not primed for a response. By pressure infiltrating the bacteria into the plant leaf, the stomata barrier was by-passed and greater infection resulted.

This simple model ignores two important facts though: the ability of stomata to close upon perception of bacteria [154] and that some bacteria synthesise and secrete coronatine enabling them to re-open closed stomata [154]. A close look at the lifecycle of *P. syringae* proved insightful in understanding how it infects plants.

Studying the life cycle of *P. syringae* indicates that it perhaps infects with greatest success during the early morning period. *P. syringae* occurs in the phyllosphere of plants and microscopy work shows that it is found in aggregates of 100 cells or more on leaf surfaces [186]. Leaf surfaces are nutrient poor areas, akin to a desert with oases of nutrients [187,188,189] and bacteria aggregates have been shown to cluster at the nutrient rich sites. Swarming is a key mechanism used by individual cells to rapidly migrate across the leaf surface in search of these nutrient rich oases [190]. Interestingly, when aggregates form at nutrient rich sites one of the early effects is repression of motility by quorum sensing [186,190,191]. It is advantageous for the small colony to stay at the nutrient rich 'oasis' rather than move away from it into the desert surroundings [186]. The plant leaf surface is a hostile area and in addition to scarce nutrients, strong sunlight and UV irradiation occurs along with exposure to free radicals such as H₂O₂ an effect of photorespiration [148,188,189]. One of the mechanisms bacteria have evolved to deal with this is stress avoidance [189]. UV and sunlight irradiation causes DNA damage possibly favouring bacterial replication at night [192]. Bacterial aggregates at nutrient rich sites replicate to populations reaching 10⁸ cells/g of leaf [186]. It has been reported that larger bacterial colonies show greater motility than smaller colonies [191] suggesting as colony size increases a second quorum is reached activating virulence genes such as coronatine production and restoring motility [191].

Thus, if aggregates of 100 cells or more cluster at nutrient rich sites replicating mainly at night, the cell population will be at late log phase, approaching stationary phase at dawn, when virulence genes are switched on by quorum sensing leading to restored motility and production of phytotoxins such as coronatine [191,193]. At dawn plant stomata open under circadian regulation [153] coinciding with peak bacterial population and greatest risk of pathogen entry. Quite possibly, in addition to anticipating *Hpa* spores, plants also anticipate bacterial invasion, thus expressing a pulse in defence gene expression at dawn. These observations could help support the model proposed here, however it will be necessary to carry out elaborate field experiments attempting to mimic natural infection. Identifying mutations in plants with constitutively open stomata would give an indication of whether there is a direct link between up regulated defence and open stomata. The molecular underpinnings of this model could be discovered by measuring transcript changes of defence related genes in response to infection at subjective night and day using microarrays.

One of the early responses to pathogens is the alkalisation of the apoplast and an influx of Ca²⁺ [36]. This results in activation of calmodulin [194] and a number of other genes in the Ca²⁺ mediated defence pathway. Bacteria actively repress this pathway using exopolysaccharides that actively sequester any free Ca²⁺ [40]. Interestingly cellular levels of Ca²⁺ display robust circadian rhythms

with peaks and nadirs during the day and night respectively [195]. As infections were performed at subjective day (peak Ca^{2+}) and subjective night (low Ca^{2+}) one could speculate that bacteria are unable to suppress the Ca^{2+} mediated defence pathway during the day but could possibly suppress Ca^{2+} when infiltrated at night as levels could be low enough allowing bacteria to sequester any Ca^{2+} . To investigate this, Ca^{2+} levels could be measured *in vivo* using the bioluminescent Ca^{2+} -reporter aequorin, during infection at both subjective dawn and dusk under free-running conditions.

This study showed promoter activity of *CCR2* is unaffected by infection, with rhythmic expression persisting after infection under free-running conditions (Fig 6.5). However, previous reports show that the HopU1, a bacterial effector, actively targets CCR2 protein [156] resulting in attenuated defence responses. This result suggests further research should also investigate the effects of infection on protein levels of clock output pathways. As the tools for genomic and proteomic studies become accessible and affordable, a clearer picture will emerge illustrating the role of the plant circadian clock in defence responses.

Concluding remarks

There is a growing body of work illustrating that the two complex and vital networks of circadian clock regulation and defence signalling are interconnected. This report highlights the complexity and nuances presented by the plant innate immune system and offers new evidence linking clock control and defence signalling. Plants display a time-of-day variation in susceptibility to the biotrophic pathogen *P. syringae* reported here and *Hpa Emw1* [146]. This time-of-day variation is lost in arrhythmic *CCA1-ox* and *elf3-1* indicating that the circadian clock drives time-of-day variation. From data reported here, one can surmise that early defence responses perhaps play a key role in defence signalling and maybe gated by the circadian clock, giving a strong response during the day when perceived threat of infection is greater. Callose production, a classic marker of early defence showed time-of-day variation supporting this claim. Mounting evidence, some of which was reported here, shows that plants seek to find a balance between a full immune response and one that is sufficient to clear infection at minimal cost to the plant. In addition to the elegant mechanisms that were discussed earlier, data presented here suggest that circadian clock control may also play a crucial part in striking a balance in defence responses thus avoiding wastefulness. In conclusion, this thesis contributes significantly to studies understanding the link between the plant circadian clock and defence responses.

References

1. Jones JD, Dangl JL (2006) The plant immune system. *Nature* 444: 323-329.
2. Postel S, Kemmerling B (2009) Plant systems for recognition of pathogen-associated molecular patterns. *Semin Cell Dev Biol* 20: 1025-1031.
3. Zipfel C (2009) Early molecular events in PAMP-triggered immunity. *Curr Opin Plant Biol* 12: 414-420.
4. Boller T, Felix G (2009) A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol* 60: 379-406.
5. Coll NS, Epple P, Dangl JL (2011) Programmed cell death in the plant immune system. *Cell Death Differ* 18: 1247-1256.
6. Ausubel FM (2005) Are innate immune signaling pathways in plants and animals conserved? *Nat Immunol* 6: 973-979.
7. Shirasu-Hiza MM, Dionne MS, Pham LN, Ayres JS, Schneider DS (2007) Interactions between circadian rhythm and immunity in *Drosophila melanogaster*. *Curr Biol* 17: R353-355.
8. Arjona A, Sarkar DK (2006) Evidence supporting a circadian control of natural killer cell function. *Brain Behav Immun* 20: 469-476.
9. Bryant PA, Trinder J, Curtis N (2004) Sick and tired: Does sleep have a vital role in the immune system? *Nat Rev Immunol* 4: 457-467.
10. Roden LC, Ingle RA (2009) Lights, Rhythms, Infection: The Role of Light and the Circadian Clock in Determining the Outcome of Plant-Pathogen Interactions. *Plant Cell*.
11. Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner HY, et al. (1996) Systemic Acquired Resistance. *Plant Cell* 8: 1809-1819.
12. Durrant WE, Dong X (2004) Systemic acquired resistance. *Annu Rev Phytopathol* 42: 185-209.
13. Bagasra O, Prilliman KR (2004) RNA interference: the molecular immune system. *J Mol Histol* 35: 545-553.
14. Zipfel C, Felix G (2005) Plants and animals: a different taste for microbes? *Curr Opin Plant Biol* 8: 353-360.
15. Thomma BP, Nurnberger T, Joosten MH (2011) Of PAMPs and effectors: the blurred PTI-ETI dichotomy. *Plant Cell* 23: 4-15.
16. Tsuda K, Katagiri F (2010) Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. *Curr Opin Plant Biol* 13: 459-465.
17. Chinchilla D, Bauer Z, Regenass M, Boller T, Felix G (2006) The Arabidopsis receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception. *Plant Cell* 18: 465-476.
18. Kunze G, Zipfel C, Robatzek S, Niehaus K, Boller T, et al. (2004) The N terminus of bacterial elongation factor Tu elicits innate immunity in Arabidopsis plants. *Plant Cell* 16: 3496-3507.
19. Felix G, Boller T (2003) Molecular sensing of bacteria in plants. The highly conserved RNA-binding motif RNP-1 of bacterial cold shock proteins is recognized as an elicitor signal in tobacco. *J Biol Chem* 278: 6201-6208.
20. Robatzek S, Chinchilla D, Boller T (2006) Ligand-induced endocytosis of the pattern recognition receptor FLS2 in Arabidopsis. *Genes Dev* 20: 537-542.
21. Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JD, et al. (2004) Bacterial disease resistance in Arabidopsis through flagellin perception. *Nature* 428: 764-767.
22. Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JD, et al. (2006) Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts Agrobacterium-mediated transformation. *Cell* 125: 749-760.
23. Shiu SH, Blecker AB (2003) Expansion of the receptor-like kinase/Pelle gene family and receptor-like proteins in Arabidopsis. *Plant Physiol* 132: 530-543.
24. Ponstein AS, Bres-Vloemans SA, Sela-Buurlage MB, van den Elzen PJ, Melchers LS, et al. (1994) A novel pathogen- and wound-inducible tobacco (*Nicotiana tabacum*) protein with antifungal activity. *Plant Physiol* 104: 109-118.

25. Dixon RA, Harrison MJ (1990) Activation, structure, and organization of genes involved in microbial defense in plants. *Adv Genet* 28: 165-234.
26. Kauffmann S, Legrand M, Geoffroy P, Fritig B (1987) Biological function of 'pathogenesis-related' proteins: four PR proteins of tobacco have 1,3-beta-glucanase activity. *EMBO J* 6: 3209-3212.
27. Tsuda K, Sato M, Glazebrook J, Cohen JD, Katagiri F (2008) Interplay between MAMP-triggered and SA-mediated defense responses. *Plant J* 53: 763-775.
28. Tsuda K, Glazebrook J, Katagiri F (2008) The interplay between MAMP and SA signaling. *Plant Signal Behav* 3: 359-361.
29. Bestwick CS, Bennett MH, Mansfield JW (1995) Hrp Mutant of *Pseudomonas syringae* pv phaseolicola Induces Cell Wall Alterations but Not Membrane Damage Leading to the Hypersensitive Reaction in Lettuce. *Plant Physiol* 108: 503-516.
30. Gomez-Gomez L, Boller T (2000) FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis. *Mol Cell* 5: 1003-1011.
31. Jacobs AK, Lipka V, Burton RA, Panstruga R, Strizhov N, et al. (2003) An Arabidopsis Callose Synthase, GSL5, Is Required for Wound and Papillary Callose Formation. *Plant Cell* 15: 2503-2513.
32. Nishimura MT, Stein M, Hou BH, Vogel JP, Edwards H, et al. (2003) Loss of a callose synthase results in salicylic acid-dependent disease resistance. *Science* 301: 969-972.
33. Kim MG, da Cunha L, McFall AJ, Belkadir Y, DebRoy S, et al. (2005) Two *Pseudomonas syringae* type III effectors inhibit RIN4-regulated basal defense in Arabidopsis. *Cell* 121: 749-759.
34. Heese A, Hann DR, Gimenez-Ibanez S, Jones AM, He K, et al. (2007) The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proc Natl Acad Sci U S A* 104: 12217-12222.
35. Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nurnberger T, et al. (2007) A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* 448: 497-500.
36. Nicaise V, Roux M, Zipfel C (2009) Recent advances in PAMP-triggered immunity against bacteria: pattern recognition receptors watch over and raise the alarm. *Plant Physiol* 150: 1638-1647.
37. Nurnberger T, Brunner F, Kemmerling B, Piater L (2004) Innate immunity in plants and animals: striking similarities and obvious differences. *Immunol Rev* 198: 249-266.
38. Garcia-Brugger A, Lamotte O, Vandelle E, Bourque S, Lecourieux D, et al. (2006) Early signaling events induced by elicitors of plant defenses. *Mol Plant Microbe Interact* 19: 711-724.
39. Ma W, Berkowitz GA (2007) The grateful dead: calcium and cell death in plant innate immunity. *Cell Microbiol* 9: 2571-2585.
40. Aslam SN, Newman MA, Erbs G, Morrissey KL, Chinchilla D, et al. (2008) Bacterial polysaccharides suppress induced innate immunity by calcium chelation. *Curr Biol* 18: 1078-1083.
41. Meszaros T, Helfer A, Hatzimasoura E, Magyar Z, Serazetdinova L, et al. (2006) The Arabidopsis MAP kinase kinase MKK1 participates in defence responses to the bacterial elicitor flagellin. *Plant J* 48: 485-498.
42. Zhang J, Shao F, Li Y, Cui H, Chen L, et al. (2007) A *Pseudomonas syringae* effector inactivates MAPKs to suppress PAMP-induced immunity in plants. *Cell Host Microbe* 1: 175-185.
43. Petersen LN, Ingle RA, Knight MR, Denby KJ (2009) OXI1 protein kinase is required for plant immunity against *Pseudomonas syringae* in Arabidopsis. *J Exp Bot* 60: 3727-3735.
44. Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, et al. (2004) OXI1 kinase is necessary for oxidative burst-mediated signalling in Arabidopsis. *Nature* 427: 858-861.
45. Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, et al. (2002) MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature* 415: 977-983.
46. Ichimura K, Casais C, Peck SC, Shinozaki K, Shirasu K (2006) MEKK1 is required for MPK4 activation and regulates tissue-specific and temperature-dependent cell death in Arabidopsis. *J Biol Chem* 281: 36969-36976.

47. Nakagami H, Soukupova H, Schikora A, Zarsky V, Hirt H (2006) A Mitogen-activated protein kinase kinase kinase mediates reactive oxygen species homeostasis in Arabidopsis. *J Biol Chem* 281: 38697-38704.
48. Su SH, Suarez-Rodriguez MC, Krysan P (2007) Genetic interaction and phenotypic analysis of the Arabidopsis MAP kinase pathway mutations mekk1 and mpk4 suggests signaling pathway complexity. *FEBS Lett* 581: 3171-3177.
49. Suarez-Rodriguez MC, Adams-Phillips L, Liu Y, Wang H, Su SH, et al. (2007) MEKK1 is required for flg22-induced MPK4 activation in Arabidopsis plants. *Plant Physiol* 143: 661-669.
50. Schweighofer A, Kazanaviciute V, Scheikl E, Teige M, Doczi R, et al. (2007) The PP2C-type phosphatase AP2C1, which negatively regulates MPK4 and MPK6, modulates innate immunity, jasmonic acid, and ethylene levels in Arabidopsis. *Plant Cell* 19: 2213-2224.
51. Eulgem T, Somssich IE (2007) Networks of WRKY transcription factors in defense signaling. *Curr Opin Plant Biol* 10: 366-371.
52. Pandey SP, Somssich IE (2009) The role of WRKY transcription factors in plant immunity. *Plant Physiol* 150: 1648-1655.
53. Eulgem T, Rushton PJ, Robatzek S, Somssich IE (2000) The WRKY superfamily of plant transcription factors. *Trends Plant Sci* 5: 199-206.
54. Sato M, Tsuda K, Wang L, Collier J, Watanabe Y, et al. (2010) Network modeling reveals prevalent negative regulatory relationships between signaling sectors in Arabidopsis immune signaling. *PLoS Pathog* 6: e1001011.
55. Naito K, Taguchi F, Suzuki T, Inagaki Y, Toyoda K, et al. (2008) Amino acid sequence of bacterial microbe-associated molecular pattern flg22 is required for virulence. *Mol Plant Microbe Interact* 21: 1165-1174.
56. Chisholm ST, Dahlbeck D, Krishnamurthy N, Day B, Sjolander K, et al. (2005) Molecular characterization of proteolytic cleavage sites of the *Pseudomonas syringae* effector AvrRpt2. *Proc Natl Acad Sci U S A* 102: 2087-2092.
57. Liu J, Elmore JM, Coaker G (2009) Investigating the functions of the RIN4 protein complex during plant innate immune responses. *Plant Signal Behav* 4: 1107-1110.
58. Xing W, Zou Y, Liu Q, Liu J, Luo X, et al. (2007) The structural basis for activation of plant immunity by bacterial effector protein AvrPto. *Nature* 449: 243-247.
59. Xiang T, Zong N, Zou Y, Wu Y, Zhang J, et al. (2008) *Pseudomonas syringae* effector AvrPto blocks innate immunity by targeting receptor kinases. *Curr Biol* 18: 74-80.
60. Janjusevic R, Abramovitch RB, Martin GB, Stebbins CE (2006) A bacterial inhibitor of host programmed cell death defenses is an E3 ubiquitin ligase. *Science* 311: 222-226.
61. van der Hoorn RA, Kamoun S (2008) From Guard to Decoy: a new model for perception of plant pathogen effectors. *Plant Cell* 20: 2009-2017.
62. Torres MA, Jones JD, Dangl JL (2006) Reactive oxygen species signaling in response to pathogens. *Plant Physiol* 141: 373-378.
63. Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW (2003) Genome-wide analysis of NBS-LRR-encoding genes in Arabidopsis. *Plant Cell* 15: 809-834.
64. Mucyn TS, Clemente A, Andriotis VM, Balmuth AL, Oldroyd GE, et al. (2006) The tomato NBARC-LRR protein Prf interacts with Pto kinase in vivo to regulate specific plant immunity. *Plant Cell* 18: 2792-2806.
65. Ntoukakis V, Mucyn TS, Gimenez-Ibanez S, Chapman HC, Gutierrez JR, et al. (2009) Host inhibition of a bacterial virulence effector triggers immunity to infection. *Science* 324: 784-787.
66. van der Hoorn RAL, Kamoun S (2008) From Guard to Decoy: A New Model for Perception of Plant Pathogen Effectors. *The Plant Cell Online* 20: 2009-2017.
67. Zipfel C, Rathjen JP (2008) Plant immunity: AvrPto targets the frontline. *Curr Biol* 18: R218-220.

68. Mackey D, Holt BF, 3rd, Wiig A, Dangl JL (2002) RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* 108: 743-754.
69. Axtell MJ, Staskawicz BJ (2003) Initiation of RPS2-specified disease resistance in *Arabidopsis* is coupled to the AvrRpt2-directed elimination of RIN4. *Cell* 112: 369-377.
70. Medzhitov R, Janeway CA, Jr. (1997) Innate immunity: the virtues of a nonclonal system of recognition. *Cell* 91: 295-298.
71. Nurnberger T, Brunner F (2002) Innate immunity in plants and animals: emerging parallels between the recognition of general elicitors and pathogen-associated molecular patterns. *Curr Opin Plant Biol* 5: 318-324.
72. Chisholm ST, Coaker G, Day B, Staskawicz BJ (2006) Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124: 803-814.
73. Haefele DM, Lindow SE (1987) Flagellar Motility Confers Epiphytic Fitness Advantages upon *Pseudomonas syringae*. *Appl Environ Microbiol* 53: 2528-2533.
74. Kvitko BH, Park DH, Velasquez AC, Wei CF, Russell AB, et al. (2009) Deletions in the repertoire of *Pseudomonas syringae* pv. tomato DC3000 type III secretion effector genes reveal functional overlap among effectors. *PLoS Pathog* 5: e1000388.
75. Kvitko BH, Ramos AR, Morello JE, Oh HS, Collmer A (2007) Identification of harpins in *Pseudomonas syringae* pv. tomato DC3000, which are functionally similar to HrpK1 in promoting translocation of type III secretion system effectors. *J Bacteriol* 189: 8059-8072.
76. Lee J, Klusener B, Tsiamis G, Stevens C, Neyt C, et al. (2001) HrpZ(Psph) from the plant pathogen *Pseudomonas syringae* pv. phaseolicola binds to lipid bilayers and forms an ion-conducting pore in vitro. *Proc Natl Acad Sci U S A* 98: 289-294.
77. Lee SW, Han SW, Bartley LE, Ronald PC (2006) From the Academy: Colloquium review. Unique characteristics of *Xanthomonas oryzae* pv. *oryzae* AvrXa21 and implications for plant innate immunity. *Proc Natl Acad Sci U S A* 103: 18395-18400.
78. Felix G, Baureithel K, Boller T (1998) Desensitization of the perception system for chitin fragments in tomato cells. *Plant Physiol* 117: 643-650.
79. Bolton MD, van Esse HP, Vossen JH, de Jonge R, Stergiopoulos I, et al. (2008) The novel *Cladosporium fulvum* lysin motif effector Ecp6 is a virulence factor with orthologues in other fungal species. *Mol Microbiol* 69: 119-136.
80. de Jonge R, van Esse HP, Kombrink A, Shinya T, Desaki Y, et al. (2010) Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science* 329: 953-955.
81. Fradin EF, Zhang Z, Juarez Ayala JC, Castroverde CD, Nazar RN, et al. (2009) Genetic dissection of *Verticillium* wilt resistance mediated by tomato Ve1. *Plant Physiol* 150: 320-332.
82. Brunner F, Rosahl S, Lee J, Rudd JJ, Geiler C, et al. (2002) Pep-13, a plant defense-inducing pathogen-associated pattern from *Phytophthora* transglutaminases. *EMBO J* 21: 6681-6688.
83. Lee SW, Han SW, Sriyanum M, Park CJ, Seo YS, et al. (2009) A type I-secreted, sulfated peptide triggers XA21-mediated innate immunity. *Science* 326: 850-853.
84. Mishina TE, Zeier J (2007) Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in *Arabidopsis*. *Plant J* 50: 500-513.
85. Wirthmueller L, Zhang Y, Jones JD, Parker JE (2007) Nuclear accumulation of the *Arabidopsis* immune receptor RPS4 is necessary for triggering EDS1-dependent defense. *Curr Biol* 17: 2023-2029.
86. Palma K, Thorgrimsen S, Malinovsky FG, Fiil BK, Nielsen HB, et al. (2010) Autoimmunity in *Arabidopsis* acd11 is mediated by epigenetic regulation of an immune receptor. *PLoS Pathog* 6.
87. Li Y, Li S, Bi D, Cheng YT, Li X, et al. (2010) SRFR1 negatively regulates plant NB-LRR resistance protein accumulation to prevent autoimmunity. *PLoS Pathog* 6.

88. Trujillo M, Ichimura K, Casais C, Shirasu K (2008) Negative regulation of PAMP-triggered immunity by an E3 ubiquitin ligase triplet in *Arabidopsis*. *Curr Biol* 18: 1396-1401.
89. Navarro L, Zipfel C, Rowland O, Keller I, Robatzek S, et al. (2004) The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis. *Plant Physiol* 135: 1113-1128.
90. Schweigkofler W, O'Donnell K, Garbelotto M (2004) Detection and quantification of airborne conidia of *Fusarium circinatum*, the causal agent of pine pitch canker, from two California sites by using a real-time PCR approach combined with a simple spore trapping method. *Appl Environ Microbiol* 70: 3512-3520.
91. Cao H, Baldini RL, Rahme LG (2001) Common mechanisms for pathogens of plants and animals. *Annu Rev Phytopathol* 39: 259-284.
92. Iriti M, Faoro F (2007) Review of innate and specific immunity in plants and animals. *Mycopathologia* 164: 57-64.
93. Donnelly MA, Steiner TS (2002) Two nonadjacent regions in enteroaggregative *Escherichia coli* flagellin are required for activation of toll-like receptor 5. *J Biol Chem* 277: 40456-40461.
94. Keller T, Damude HG, Werner D, Doerner P, Dixon RA, et al. (1998) A plant homolog of the neutrophil NADPH oxidase gp91phox subunit gene encodes a plasma membrane protein with Ca²⁺ binding motifs. *Plant Cell* 10: 255-266.
95. Torres MA, Onouchi H, Hamada S, Machida C, Hammond-Kosack KE, et al. (1998) Six *Arabidopsis thaliana* homologues of the human respiratory burst oxidase (gp91phox). *Plant J* 14: 365-370.
96. Torres MA, Dangl JL, Jones JD (2002) *Arabidopsis* gp91phox homologues AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates in the plant defense response. *Proc Natl Acad Sci U S A* 99: 517-522.
97. Holt BF, 3rd, Hubert DA, Dangl JL (2003) Resistance gene signaling in plants--complex similarities to animal innate immunity. *Curr Opin Immunol* 15: 20-25.
98. Nimchuk Z, Eulgem T, Holt BF, 3rd, Dangl JL (2003) Recognition and response in the plant immune system. *Annu Rev Genet* 37: 579-609.
99. Edery I (2000) Circadian rhythms in a nutshell. *Physiol Genomics* 3: 59-74.
100. Harmer SL (2009) The Circadian System in Higher Plants. *Annu Rev Plant Biol*.
101. Dodd AN, Salathia N, Hall A, Kevei E, Toth R, et al. (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. *Science* 309: 630-633.
102. Hastings JW, Sweeney BM (1957) On the Mechanism of Temperature Independence in a Biological Clock. *Proc Natl Acad Sci U S A* 43: 804-811.
103. Pittendrigh CS (1993) Temporal organization: reflections of a Darwinian clock-watcher. *Annu Rev Physiol* 55: 16-54.
104. Dunlap JC (1999) Molecular bases for circadian clocks. *Cell* 96: 271-290.
105. Ashmore LJ, Sathyanarayanan S, Silvestre DW, Emerson MM, Schotland P, et al. (2003) Novel insights into the regulation of the timeless protein. *J Neurosci* 23: 7810-7819.
106. Lee C, Bae K, Edery I (1998) The *Drosophila* CLOCK protein undergoes daily rhythms in abundance, phosphorylation, and interactions with the PER-TIM complex. *Neuron* 21: 857-867.
107. Lee C, Bae K, Edery I (1999) PER and TIM inhibit the DNA binding activity of a *Drosophila* CLOCK-CYC/DBMAL1 heterodimer without disrupting formation of the heterodimer: a basis for circadian transcription. *Mol Cell Biol* 19: 5316-5325.
108. Darlington TK, Wager-Smith K, Ceriani MF, Staknis D, Gekakis N, et al. (1998) Closing the circadian loop: CLOCK-induced transcription of its own inhibitors per and tim. *Science* 280: 1599-1603.
109. So WV, Rosbash M (1997) Post-transcriptional regulation contributes to *Drosophila* clock gene mRNA cycling. *EMBO J* 16: 7146-7155.

110. Zeng H, Hardin PE, Rosbash M (1994) Constitutive overexpression of the *Drosophila* period protein inhibits period mRNA cycling. *EMBO J* 13: 3590-3598.
111. Price JL, Blau J, Rothenfluh A, Abodeely M, Kloss B, et al. (1998) double-time is a novel *Drosophila* clock gene that regulates PERIOD protein accumulation. *Cell* 94: 83-95.
112. Naidoo N, Song W, Hunter-Ensor M, Sehgal A (1999) A role for the proteasome in the light response of the timeless clock protein. *Science* 285: 1737-1741.
113. Kloss B, Price JL, Saez L, Blau J, Rothenfluh A, et al. (1998) The *Drosophila* clock gene double-time encodes a protein closely related to human casein kinase Iε. *Cell* 94: 97-107.
114. Vossell LB, Price JL, Sehgal A, Saez L, Young MW (1994) Block in nuclear localization of period protein by a second clock mutation, timeless. *Science* 263: 1606-1609.
115. Myers MP, Wager-Smith K, Rothenfluh-Hilfiker A, Young MW (1996) Light-induced degradation of TIMELESS and entrainment of the *Drosophila* circadian clock. *Science* 271: 1736-1740.
116. Saez L, Young MW (1996) Regulation of nuclear entry of the *Drosophila* clock proteins period and timeless. *Neuron* 17: 911-920.
117. Edery I, Zwiebel LJ, Dembinska ME, Rosbash M (1994) Temporal phosphorylation of the *Drosophila* period protein. *Proc Natl Acad Sci U S A* 91: 2260-2264.
118. Bae K, Lee C, Sidote D, Chuang KY, Edery I (1998) Circadian regulation of a *Drosophila* homolog of the mammalian Clock gene: PER and TIM function as positive regulators. *Mol Cell Biol* 18: 6142-6151.
119. Lee JE, Edery I (2008) Circadian regulation in the ability of *Drosophila* to combat pathogenic infections. *Curr Biol* 18: 195-199.
120. McDonald MJ, Rosbash M (2001) Microarray analysis and organization of circadian gene expression in *Drosophila*. *Cell* 107: 567-578.
121. Bischoff V, Vignal C, Duvic B, Boneca IG, Hoffmann JA, et al. (2006) Downregulation of the *Drosophila* immune response by peptidoglycan-recognition proteins SC1 and SC2. *PLoS Pathog* 2: e14.
122. Mukae N, Yokoyama H, Yokokura T, Sakoyama Y, Nagata S (2002) Activation of the innate immunity in *Drosophila* by endogenous chromosomal DNA that escaped apoptotic degradation. *Genes Dev* 16: 2662-2671.
123. Tsuda M, Langmann C, Harden N, Aigaki T (2005) The RING-finger scaffold protein Plenty of SH3s targets TAK1 to control immunity signalling in *Drosophila*. *EMBO Rep* 6: 1082-1087.
124. Reppert SM, Weaver DR (2002) Coordination of circadian timing in mammals. *Nature* 418: 935-941.
125. Arjona A, Boyadjieva N, Sarkar DK (2004) Circadian rhythms of granzyme B, perforin, IFN-γ, and NK cell cytolytic activity in the spleen: effects of chronic ethanol. *J Immunol* 172: 2811-2817.
126. Buguet A, Bert J, Tapie P, Tabaraud F, Doua F, et al. (1993) Sleep-wake cycle in human African trypanosomiasis. *J Clin Neurophysiol* 10: 190-196.
127. Born J, Lange T, Hansen K, Molle M, Fehm HL (1997) Effects of sleep and circadian rhythm on human circulating immune cells. *J Immunol* 158: 4454-4464.
128. Heiser P, Dickhaus B, Schreiber W, Clement HW, Hasse C, et al. (2000) White blood cells and cortisol after sleep deprivation and recovery sleep in humans. *Eur Arch Psychiatry Clin Neurosci* 250: 16-23.
129. Yakir E, Hassidim M, Melamed-Book N, Hilman D, Kron I, et al. (2011) Cell Autonomous and Cell-Type Specific Circadian Rhythms in *Arabidopsis*. *Plant J*.
130. Jones MA, Covington MF, DiTacchio L, Vollmers C, Panda S, et al. (2010) Jumonji domain protein JMJD5 functions in both the plant and human circadian systems. *Proc Natl Acad Sci U S A* 107: 21623-21628.
131. McClung CR (2009) Circadian rhythms. Linking the loops. *Science* 323: 1440-1441.
132. Alabadi D, Oyama T, Yanovsky MJ, Harmon FG, Mas P, et al. (2001) Reciprocal regulation between TOC1 and LHY/CCA1 within the *Arabidopsis* circadian clock. *Science* 293: 880-883.

133. Pruneda-Paz JL, Breton G, Para A, Kay SA (2009) A Functional Genomics Approach Reveals CHE as a Component of the Arabidopsis Circadian Clock. *Science* 323: 1481-1485.
134. Yakir E, Hilman D, Kron I, Hassidim M, Melamed-Book N, et al. (2009) Posttranslational regulation of CIRCADIAN CLOCK ASSOCIATED1 in the circadian oscillator of Arabidopsis. *Plant Physiol* 150: 844-857.
135. Lu SX, Knowles SM, Andronis C, Ong MS, Tobin EM (2009) CIRCADIAN CLOCK ASSOCIATED1 and LATE ELONGATED HYPOCOTYL function synergistically in the circadian clock of Arabidopsis. *Plant Physiol* 150: 834-843.
136. Streitner C, Hennig L, Korneli C, Staiger D (2010) Global transcript profiling of transgenic plants constitutively overexpressing the RNA-binding protein AtGRP7. *BMC Plant Biol* 10: 221.
137. Covington MF, Maloof JN, Straume M, Kay SA, Harmer SL (2008) Global transcriptome analysis reveals circadian regulation of key pathways in plant growth and development. *Genome Biol* 9: R130.
138. Michael TP, McClung CR (2003) Enhancer trapping reveals widespread circadian clock transcriptional control in Arabidopsis. *Plant Physiol* 132: 629-639.
139. Edwards KD, Anderson PE, Hall A, Salathia NS, Locke JC, et al. (2006) FLOWERING LOCUS C mediates natural variation in the high-temperature response of the Arabidopsis circadian clock. *Plant Cell* 18: 639-650.
140. Harmer SL, Hogenesch JB, Straume M, Chang HS, Han B, et al. (2000) Orchestrated transcription of key pathways in Arabidopsis by the circadian clock. *Science* 290: 2110-2113.
141. Harmer SL, Kay SA (2005) Positive and negative factors confer phase-specific circadian regulation of transcription in Arabidopsis. *Plant Cell* 17: 1926-1940.
142. Michael TP, Mockler TC, Breton G, McEntee C, Byer A, et al. (2008) Network discovery pipeline elucidates conserved time-of-day-specific cis-regulatory modules. *PLoS Genet* 4: e14.
143. Hudson ME, Quail PH (2003) Identification of promoter motifs involved in the network of phytochrome A-regulated gene expression by combined analysis of genomic sequence and microarray data. *Plant Physiol* 133: 1605-1616.
144. Salter MG, Franklin KA, Whitelam GC (2003) Gating of the rapid shade-avoidance response by the circadian clock in plants. *Nature* 426: 680-683.
145. Franklin KA, Praekelt U, Stoddart WM, Billingham OE, Halliday KJ, et al. (2003) Phytochromes B, D, and E act redundantly to control multiple physiological responses in Arabidopsis. *Plant Physiol* 131: 1340-1346.
146. Wang W, Barnaby JY, Tada Y, Li H, Tor M, et al. (2011) Timing of plant immune responses by a central circadian regulator. *Nature* 470: 110-114.
147. Dodd RS, Huberli D, Mayer W, Harnik TY, Afzal-Rafli Z, et al. (2008) Evidence for the role of synchronicity between host phenology and pathogen activity in the distribution of sudden oak death canker disease. *New Phytol* 179: 505-514.
148. Hirano SS, Upper CD (1991) Bacterial community dynamics in Microbial Ecology of Leaves. In: Andrews JHaH, S.S., editor. NY: New York- Springer-Verlag. pp. 271-294.
149. Weyman PD, Pan Z, Feng Q, Gilchrist DG, Bostock RM (2006) A circadian rhythm-regulated tomato gene is induced by Arachidonic acid and *Phytophthora infestans* infection. *Plant Physiol* 140: 235-248.
150. Sauerbrunn N, Schlaich NL (2004) PCC1: a merging point for pathogen defence and circadian signalling in Arabidopsis. *Planta* 218: 552-561.
151. Faris JD, Zhang Z, Lu H, Lu S, Reddy L, et al. (2010) A unique wheat disease resistance-like gene governs effector-triggered susceptibility to necrotrophic pathogens. *Proc Natl Acad Sci U S A* 107: 13544-13549.
152. Molina A, Mena M, Carbonero P, Garcia-Olmedo F (1997) Differential expression of pathogen-responsive genes encoding two types of glycine-rich proteins in barley. *Plant Mol Biol* 33: 803-810.

153. Somers DE, Webb AA, Pearson M, Kay SA (1998) The short-period mutant, *toc1-1*, alters circadian clock regulation of multiple outputs throughout development in *Arabidopsis thaliana*. *Development* 125: 485-494.
154. Melotto M, Underwood W, Koczan J, Nomura K, He SY (2006) Plant stomata function in innate immunity against bacterial invasion. *Cell* 126: 969-980.
155. Kim JS, Jung HJ, Lee HJ, Kim KA, Goh CH, et al. (2008) Glycine-rich RNA-binding protein 7 affects abiotic stress responses by regulating stomata opening and closing in *Arabidopsis thaliana*. *Plant J* 55: 455-466.
156. Fu ZQ, Guo M, Jeong BR, Tian F, Elthon TE, et al. (2007) A type III effector ADP-ribosylates RNA-binding proteins and quells plant immunity. *Nature* 447: 284-288.
157. Burgess A, Searle I (2011) The clock primes defense at dawn. *Immunol Cell Biol*.
158. Murray SL, Thomson C, Chini A, Read ND, Loake GJ (2002) Characterization of a novel, defense-related *Arabidopsis* mutant, *cir1*, isolated by luciferase imaging. *Mol Plant Microbe Interact* 15: 557-566.
159. King EO, Ward MK, Raney DE (1954) Two simple media for the demonstration of pyocyanin and fluorescein. *J Lab Clin Med* 44: 301-307.
160. Katagiri F, Thilmony, R. & He, S.Y. , editor (2002) *The Arabidopsis thaliana-Pseudomonas syringae* Interaction. Rockville: Am. Soc. Plant Biologists, Rockville, MD. 1-35 p.
161. Ewbank J, Vivier E, Kim MG, Mackey D (2008) Measuring Cell-Wall-Based Defenses and Their Effect on Bacterial Growth in *Arabidopsis*. *Innate Immunity*: Humana Press. pp. 443-452.
162. Smart M, Roden LC (2010) A small-scale RNA isolation protocol useful for high-throughput extractions from recalcitrant plants. *South African Journal of Botany* 76: 375-379.
163. Niinuma K, Someya N, Kimura M, Yamaguchi I, Hamamoto H (2005) Circadian rhythm of circumnutation in inflorescence stems of *Arabidopsis*. *Plant Cell Physiol* 46: 1423-1427.
164. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29: e45.
165. Rasmussen R (2001) Quantification on the LightCycler.; Meuer S, Wittwer, C. and Nakagawara, KK. (eds), editor. Heidelberg: Springer Press. 21-34 p.
166. Wang ZY, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. *Cell* 93: 1207-1217.
167. Zhou X, Jiang Y, Yu D (2011) WRKY22 transcription factor mediates dark-induced leaf senescence in *Arabidopsis*. *Mol Cells* 31: 303-313.
168. Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, et al. (2008) Activation of defense response pathways by OGs and Flg22 elicitors in *Arabidopsis* seedlings. *Mol Plant* 1: 423-445.
169. Kwon SI, Kim SH, Bhattacharjee S, Noh JJ, Gassmann W (2009) SRFR1, a suppressor of effector-triggered immunity, encodes a conserved tetratricopeptide repeat protein with similarity to transcriptional repressors. *Plant J* 57: 109-119.
170. Hicks KA, Millar AJ, Carre IA, Somers DE, Straume M, et al. (1996) Conditional circadian dysfunction of the *Arabidopsis* early-flowering 3 mutant. *Science* 274: 790-792.
171. Cui H, Wang Y, Xue L, Chu J, Yan C, et al. (2010) *Pseudomonas syringae* effector protein AvrB perturbs *Arabidopsis* hormone signaling by activating MAP kinase 4. *Cell Host Microbe* 7: 164-175.
172. Innes RW, Bent AF, Kunkel BN, Bisgrove SR, Staskawicz BJ (1993) Molecular analysis of avirulence gene *avrRpt2* and identification of a putative regulatory sequence common to all known *Pseudomonas syringae* avirulence genes. *J Bacteriol* 175: 4859-4869.
173. Dong J, Chen C, Chen Z (2003) Expression profiles of the *Arabidopsis* WRKY gene superfamily during plant defense response. *Plant Mol Biol* 51: 21-37.

174. Rogers EE, Ausubel FM (1997) Arabidopsis enhanced disease susceptibility mutants exhibit enhanced susceptibility to several bacterial pathogens and alterations in PR-1 gene expression. *Plant Cell* 9: 305-316.
175. Friedrich L, Lawton K, Dietrich R, Willits M, Cade R, et al. (2001) NIM1 overexpression in Arabidopsis potentiates plant disease resistance and results in enhanced effectiveness of fungicides. *Mol Plant Microbe Interact* 14: 1114-1124.
176. Kim SH, Gao F, Bhattacharjee S, Adiasor JA, Nam JC, et al. (2010) The Arabidopsis resistance-like gene SNC1 is activated by mutations in SRFR1 and contributes to resistance to the bacterial effector AvrRps4. *PLoS Pathog* 6: e1001172.
177. Kim SH, Kwon SI, Saha D, Anyanwu NC, Gassmann W (2009) Resistance to the *Pseudomonas syringae* effector HopA1 is governed by the TIR-NBS-LRR protein RPS6 and is enhanced by mutations in SRFR1. *Plant Physiol* 150: 1723-1732.
178. Hall A, Bastow RM, Davis SJ, Hanano S, McWatters HG, et al. (2003) The TIME FOR COFFEE gene maintains the amplitude and timing of Arabidopsis circadian clocks. *Plant Cell* 15: 2719-2729.
179. Carpenter CD, Kreps JA, Simon AE (1994) Genes encoding glycine-rich Arabidopsis thaliana proteins with RNA-binding motifs are influenced by cold treatment and an endogenous circadian rhythm. *Plant Physiol* 104: 1015-1025.
180. Heintzen C, Melzer S, Fischer R, Kappeler S, Apel K, et al. (1994) A light- and temperature-entrained circadian clock controls expression of transcripts encoding nuclear proteins with homology to RNA-binding proteins in meristematic tissue. *Plant J* 5: 799-813.
181. Heintzen C, Nater M, Apel K, Staiger D (1997) AtGRP7, a nuclear RNA-binding protein as a component of a circadian-regulated negative feedback loop in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* 94: 8515-8520.
182. Griebel T, Zeier J (2008) Light regulation and daytime dependency of inducible plant defenses in Arabidopsis: phytochrome signaling controls systemic acquired resistance rather than local defense. *Plant Physiol* 147: 790-801.
183. Green RM, Tingay S, Wang ZY, Tobin EM (2002) Circadian rhythms confer a higher level of fitness to Arabidopsis plants. *Plant Physiol* 129: 576-584.
184. McClung CR (2011) Plant biology: Defence at dawn. *Nature* 470: 44-45.
185. Donofrio NM, Delaney TP (2001) Abnormal callose response phenotype and hypersusceptibility to *Peronospora parasitica* in defence-compromised arabidopsis nim1-1 and salicylate hydroxylase-expressing plants. *Mol Plant Microbe Interact* 14: 439-450.
186. Monier JM, Lindow SE (2003) Differential survival of solitary and aggregated bacterial cells promotes aggregate formation on leaf surfaces. *Proc Natl Acad Sci U S A* 100: 15977-15982.
187. Leveau JH, Lindow SE (2001) Appetite of an epiphyte: quantitative monitoring of bacterial sugar consumption in the phyllosphere. *Proc Natl Acad Sci U S A* 98: 3446-3453.
188. Hirano SS, Upper CD (2000) Bacteria in the leaf ecosystem with emphasis on *Pseudomonas syringae*-a pathogen, ice nucleus, and epiphyte. *Microbiol Mol Biol Rev* 64: 624-653.
189. Wilson M, Hirano SS, Lindow SE (1999) Location and survival of leaf-associated bacteria in relation to pathogenicity and potential for growth within the leaf. *Appl Environ Microbiol* 65: 1435-1443.
190. Kinscherf TG, Willis DK (1999) Swarming by *Pseudomonas syringae* B728a requires gacS (IemA) and gacA but not the acyl-homoserine lactone biosynthetic gene ahII. *J Bacteriol* 181: 4133-4136.
191. Quinones B, Dulla G, Lindow SE (2005) Quorum sensing regulates exopolysaccharide production, motility, and virulence in *Pseudomonas syringae*. *Mol Plant Microbe Interact* 18: 682-693.
192. Iriarte FB, Balogh B, Momol MT, Smith LM, Wilson M, et al. (2007) Factors affecting survival of bacteriophage on tomato leaf surfaces. *Appl Environ Microbiol* 73: 1704-1711.

193. Liu H, Coulthurst SJ, Pritchard L, Hedley PE, Ravensdale M, et al. (2008) Quorum sensing coordinates brute force and stealth modes of infection in the plant pathogen *Pectobacterium atrosepticum*. PLoS Pathog 4: e1000093.
194. Reddy VS, Reddy AS (2004) Proteomics of calcium-signaling components in plants. Phytochemistry 65: 1745-1776.
195. Love J, Dodd AN, Webb AA (2004) Circadian and diurnal calcium oscillations encode photoperiodic information in *Arabidopsis*. Plant Cell 16: 956-966.

University of Cape Town